


## Anti-Hip antibody



<b>Description</b>	Mouse monoclonal to Hip.
<b>Model</b>	STJ98136
<b>Host</b>	Mouse
<b>Reactivity</b>	Human, Simian
<b>Applications</b>	ELISA, IF, IHC, WB
<b>Immunogen</b>	Purified recombinant fragment of human Hip expressed in E. Coli.
<b>Gene ID</b>	<a href="#">6767</a>
<b>Gene Symbol</b>	<a href="#">ST13</a>
<b>Dilution range</b>	WB 1:500-1:2000IHC 1:200-1:1000IF 1:200-1:1000ELISA 1:10000
<b>Specificity</b>	Hip Monoclonal Antibody detects endogenous levels of Hip protein.
<b>Purification</b>	Affinity purification
<b>Clone ID</b>	3B10
<b>Note</b>	For Research Use Only (RUO).
<b>Protein Name</b>	Hsc70-interacting protein Hip Aging-associated protein 2 Progesterone receptor-associated p48 protein Protein FAM10A1 Putative tumor suppressor ST13 Renal carcinoma antigen NY-REN-33 Suppression of tumorigenicity
<b>Clonality</b>	Monoclonal
<b>Conjugation</b>	Unconjugated
<b>Isotype</b>	IgG1

<b>Formulation</b>	Ascitic fluid containing 0.03% sodium azide.
<b>Storage Instruction</b>	Store at -20°C, and avoid repeat freeze-thaw cycles.
<b>Database Links</b>	<a href="#">HGNC:11343</a> <a href="#">OMIM:606796</a>
<b>Alternative Names</b>	Hsc70-interacting protein Hip Aging-associated protein 2 Progesterone receptor-associated p48 protein Protein FAM10A1 Putative tumor suppressor ST13 Renal carcinoma antigen NY-REN-33 Suppression of tumorigenicity
<b>Function</b>	One HIP oligomer binds the ATPase domains of at least two HSC70 molecules dependent on activation of the HSC70 ATPase by HSP40. Stabilizes the ADP state of HSC70 that has a high affinity for substrate protein. Through its own chaperone activity, it may contribute to the interaction of HSC70 with various target proteins .
<b>Cellular Localization</b>	Cytoplasm

---

**St John's Laboratory Ltd**

**F** +44 (0)207 681 2580  
**T** +44 (0)208 223 3081

**W** <http://www.stjohnslabs.com/>  
**E** [info@stjohnslabs.com](mailto:info@stjohnslabs.com)