


Anti-HSP60 antibody



Description Mouse monoclonal to HSP60.

Model STJ98142

Host Mouse

Reactivity Human, Mouse, Rat, Simian

Applications ELISA, FC, IF, IHC, WB

Immunogen Purified recombinant fragment of human HSP60 expressed in E. Coli.

Gene ID 3329

Gene Symbol HSPD1

Dilution range WB 1:500-1:2000IHC 1:200-1:1000IF 1:200-1:1000FC 1:200-1:400ELISA

1:10000

Specificity HSP60 Monoclonal Antibody detects endogenous levels of HSP60 protein.

Purification Affinity purification

Clone ID 3G8

Note For Research Use Only (RUO).

Protein Name 60 kDa heat shock protein, mitochondrial 60 kDa chaperonin Chaperonin 60

CPN60 Heat shock protein 60 HSP-60 Hsp60 HuCHA60 Mitochondrial

matrix protein P1 P60 lymphocyte protein

Clonality Monoclonal

Conjugation Unconjugated

Isotype IgG1

Formulation Ascitic fluid containing 0.03% sodium azide.

Store at -20°C, and avoid repeat freeze-thaw cycles. **Storage Instruction**

HGNC:52610MIM:118190 **Database Links**

60 kDa heat shock protein, mitochondrial 60 kDa chaperonin Chaperonin 60 **Alternative Names**

CPN60 Heat shock protein 60 HSP-60 Hsp60 HuCHA60 Mitochondrial

matrix protein P1 P60 lymphocyte protein

Function Chaperonin implicated in mitochondrial protein import and macromolecular

> assembly. Together with Hsp10, facilitates the correct folding of imported proteins. May also prevent misfolding and promote the refolding and proper assembly of unfolded polypeptides generated under stress conditions in the mitochondrial matrix. The functional units of these chaperonins consist of heptameric rings of the large subunit Hsp60, which function as a back-to-back double ring. In a cyclic reaction, Hsp60 ring complexes bind one unfolded substrate protein per ring, followed by the binding of ATP and association with 2 heptameric rings of the co-chaperonin Hsp10. This leads to sequestration of the substrate protein in the inner cavity of Hsp60 where, for a certain period of time, it can fold undisturbed by other cell components. Synchronous hydrolysis of ATP in all Hsp60 subunits results in the

dissociation of the chaperonin rings and the release of ADP and the folded substrate protein (Probable).

Cellular Localization Mitochondrion matrix.

St John's Laboratory Ltd

F +44 (0)207 681 2580

W http://www.stjohnslabs.com/ T +44 (0)208 223 3081 E info@stjohnslabs.com