


Anti-Ret antibody



Description	Mouse monoclonal to Ret.
Model	STJ98358
Host	Mouse
Reactivity	Human
Applications	ELISA, WB
Immunogen	Purified recombinant fragment of Ret (aa896-1063) expressed in E. Coli.
Immunogen Region	896-1063aa
Gene ID	5979
Gene Symbol	RET
Dilution range	WB 1:500-1:2000ELISA 1:10000
Specificity	Ret Monoclonal Antibody detects endogenous levels of Ret protein.
Purification	Affinity purification
Clone ID	8D10C9
Note	For Research Use Only (RUO).
Protein Name	Proto-oncogene tyrosine-protein kinase receptor Ret Cadherin family member 12 Proto-oncogene c-Ret Soluble RET kinase fragment Extracellular cell-membrane anchored RET cadherin 120 kDa fragment
Clonality	Monoclonal
Conjugation	Unconjugated

Isotype	IgG1
Formulation	Ascitic fluid containing 0.03% sodium azide.
Storage Instruction	Store at -20°C, and avoid repeat freeze-thaw cycles.
Database Links	HGNC:9967OMIM:114500
Alternative Names	Proto-oncogene tyrosine-protein kinase receptor Ret Cadherin family member 12 Proto-oncogene c-Ret Soluble RET kinase fragment Extracellular cell-membrane anchored RET cadherin 120 kDa fragment
Function	Receptor tyrosine-protein kinase involved in numerous cellular mechanisms including cell proliferation, neuronal navigation, cell migration, and cell differentiation upon binding with glial cell derived neurotrophic factor family ligands. Phosphorylates PTK2/FAK1. Regulates both cell death/survival balance and positional information. Required for the molecular mechanisms orchestration during intestine organogenesis; involved in the development of enteric nervous system and renal organogenesis during embryonic life, and promotes the formation of Peyer's patch-like structures, a major component of the gut-associated lymphoid tissue. Modulates cell adhesion via its cleavage by caspase in sympathetic neurons and mediates cell migration in an integrin (e.g. ITGB1 and ITGB3)-dependent manner. Involved in the development of the neural crest. Active in the absence of ligand, triggering apoptosis through a mechanism that requires receptor intracellular caspase cleavage. Acts as a dependence receptor; in the presence of the ligand GDNF in somatotrophs (within pituitary), promotes survival and down regulates growth hormone (GH) production, but triggers apoptosis in absence of GDNF. Regulates nociceptor survival and size. Triggers the differentiation of rapidly adapting (RA) mechanoreceptors. Mediator of several diseases such as neuroendocrine cancers; these diseases are characterized by aberrant integrins-regulated cell migration.
Cellular Localization	Cell membrane Endosome membrane
Post-translational Modifications	Autophosphorylated on C-terminal tyrosine residues upon ligand stimulation. Dephosphorylated by PTPRJ on Tyr-905, Tyr-1015 and Tyr-1062. Proteolytically cleaved by caspase-3. The soluble RET kinase fragment is able to induce cell death. The extracellular cell-membrane anchored RET cadherin fragment accelerates cell adhesion in sympathetic neurons.