


Anti-Wnt-1 antibody



Description	Mouse monoclonal to Wnt-1.
Model	STJ98451
Host	Mouse
Reactivity	Human, Mouse
Applications	ELISA, FC, IF, IHC, WB
Immunogen	Purified recombinant fragment of Wnt-1 expressed in E. Coli.
Gene ID	7471
Gene Symbol	WNT1
Dilution range	WB 1:500-1:2000IHC 1:200-1:1000IF 1:200-1:1000FC 1:200-1:400ELISA 1:10000
Specificity	Wnt-1 Monoclonal Antibody detects endogenous levels of Wnt-1 protein.
Purification	Affinity purification
Clone ID	10C8
Note	For Research Use Only (RUO).
Protein Name	Proto-oncogene Wnt-1 Proto-oncogene Int-1 homolog
Clonality	Monoclonal
Conjugation	Unconjugated
Isotype	IgG1
Formulation	Ascitic fluid containing 0.03% sodium azide.

Storage Instruction	Store at -20°C, and avoid repeat freeze-thaw cycles.
Database Links	HGNC:12774OMIM:164820
Alternative Names	Proto-oncogene Wnt-1 Proto-oncogene Int-1 homolog
Function	Ligand for members of the frizzled family of seven transmembrane receptors. In some developmental processes, is also a ligand for the coreceptor RYK, thus triggering Wnt signaling. Probable developmental protein. May be a signaling molecule important in CNS development. Is likely to signal over only few cell diameters. Has a role in osteoblast function and bone development.
Cellular Localization	Secreted, extracellular space, extracellular matrix.
Post-translational Modifications	Palmitoleylation is required for efficient binding to frizzled receptors. Palmitoleylation is necessary for proper trafficking to cell surface (Probable). Depalmitoleylated by NOTUM, leading to inhibit Wnt signaling pathway .

St John's Laboratory Ltd

F +44 (0)207 681 2580

T +44 (0)208 223 3081

W <http://www.stjohnslabs.com/>

E info@stjohnslabs.com