


Anti-TFIIH p62 antibody



Description	Mouse monoclonal to TFIIH p62.
Model	STJ98561
Host	Mouse
Reactivity	Bovine, Canine, Human, Mouse, Rat, Swine
Applications	WB
Immunogen	Purified recombinant human TFIIH p62 (N-terminal) protein fragments expressed in E.coli.
Immunogen Region	N-terminal
Gene ID	2965
Gene Symbol	GTF2H1
Dilution range	WB 1:1000-1:2000
Specificity	TFIIH p62 Monoclonal Antibody detects endogenous levels of TFIIH p62 protein.
Purification	Affinity purification
Note	For Research Use Only (RUO).
Protein Name	General transcription factor IIH subunit 1 Basic transcription factor 2 62 kDa subunit BTF2 p62 General transcription factor IIH polypeptide 1 TFIIH basal transcription factor complex p62 subunit
Clonality	Monoclonal
Conjugation	Unconjugated

Formulation	Purified mouse monoclonal in buffer containing 0.1M Tris-Glycine (pH 7.4, 150 mM NaCl) with 0.2% sodium azide, 50% glycerol.
Concentration	1 mg/ml
Storage Instruction	Store at -20°C, and avoid repeat freeze-thaw cycles.
Database Links	HGNC:4655OMIM:189972
Alternative Names	General transcription factor IIH subunit 1 Basic transcription factor 2 62 kDa subunit BTF2 p62 General transcription factor IIH polypeptide 1 TFIID basal transcription factor complex p62 subunit
Function	Component of the core-TFIID basal transcription factor involved in nucleotide excision repair (NER) of DNA and, when complexed to CAK, in RNA transcription by RNA polymerase II.
Cellular Localization	Nucleus.

St John's Laboratory Ltd

F +44 (0)207 681 2580
T +44 (0)208 223 3081

W <http://www.stjohnslabs.com/>
E info@stjohnslabs.com