


## Anti-DDR2 antibody



<b>Description</b>	Rabbit polyclonal to DDR2.
<b>Model</b>	STJ98753
<b>Host</b>	Rabbit
<b>Reactivity</b>	Human, Mouse
<b>Applications</b>	ELISA, IHC
<b>Immunogen</b>	Synthetic peptide from human DDR2 protein.
<b>Immunogen Region</b>	31-80 aa
<b>Gene ID</b>	<a href="#">4921</a>
<b>Gene Symbol</b>	<a href="#">DDR2</a>
<b>Dilution range</b>	IHC-P 1:50-300ELISA 1:5000-20000
<b>Specificity</b>	The antibody detects endogenous DDR2.
<b>Tissue Specificity</b>	Detected in osteocytes, osteoblastic cells in subchondral bone, bone lining cells, tibia and cartilage (at protein level). Detected at high levels in heart and lung, and at low levels in brain, placenta, liver, skeletal muscle, pancreas, and kidney.
<b>Purification</b>	The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen.
<b>Note</b>	For Research Use Only (RUO).
<b>Protein Name</b>	Discoidin domain-containing receptor 2 Discoidin domain receptor 2 CD167 antigen-like family member B Discoidin domain-containing receptor tyrosine kinase 2 Neurotrophic tyrosine kinase, receptor-related 3 Receptor protein-ty

<b>Clonality</b>	Polyclonal
<b>Conjugation</b>	Unconjugated
<b>Isotype</b>	IgG
<b>Formulation</b>	PBS, pH 7.4, containing 0.02% sodium azide as Preservative and 50% Glycerol.
<b>Concentration</b>	1 mg/ml
<b>Storage Instruction</b>	Store at -20°C, and avoid repeat freeze-thaw cycles.
<b>Database Links</b>	<a href="#">HGNC:2731OMIM:191311</a>
<b>Alternative Names</b>	Discoidin domain-containing receptor 2 Discoidin domain receptor 2 CD167 antigen-like family member B Discoidin domain-containing receptor tyrosine kinase 2 Neurotrophic tyrosine kinase, receptor-related 3 Receptor protein-ty
<b>Function</b>	Tyrosine kinase that functions as cell surface receptor for fibrillar collagen and regulates cell differentiation, remodeling of the extracellular matrix, cell migration and cell proliferation. Required for normal bone development. Regulates osteoblast differentiation and chondrocyte maturation via a signaling pathway that involves MAP kinases and leads to the activation of the transcription factor RUNX2. Regulates remodeling of the extracellular matrix by up-regulation of the collagenases MMP1, MMP2 and MMP13, and thereby facilitates cell migration and tumor cell invasion. Promotes fibroblast migration and proliferation, and thereby contributes to cutaneous wound healing.
<b>Cellular Localization</b>	Cell membrane
<b>Post-translational Modifications</b>	N-glycosylated. Tyrosine phosphorylated in response to collagen binding. Phosphorylated by SRC; this is required for activation and subsequent autophosphorylation on additional tyrosine residues.