


Anti-TG beta antibody



Description	Rabbit polyclonal to TGFbeta2.
Model	STJ98769
Host	Rabbit
Reactivity	Human, Mouse, Rat
Applications	ELISA, IHC, WB
Immunogen	Synthetic peptide from human TGFbeta2 protein.
Immunogen Region	61-110 aa
Gene ID	7042
Gene Symbol	TGFB2
Dilution range	WB 1:500-2000IHC-P 1:50-300ELISA 1:5000-20000
Specificity	The antibody detects endogenous TGFbeta2.
Purification	The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen.
Note	For Research Use Only (RUO).
Protein Name	Transforming growth factor beta-2 TGF-beta-2 BSC-1 cell growth inhibitor Cetermin Glioblastoma-derived T-cell suppressor factor G-TSF Polyergin Latency-associated peptide LAP
Molecular Weight	47+12kDa
Clonality	Polyclonal
Conjugation	Unconjugated

Isotype	IgG
Formulation	PBS, pH 7.4, containing 0.02% sodium azide as Preservative and 50% Glycerol.
Concentration	1 mg/ml
Storage Instruction	Store at -20°C, and avoid repeat freeze-thaw cycles.
Database Links	HGNC:11768OMIM:190220
Alternative Names	Transforming growth factor beta-2 TGF-beta-2 BSC-1 cell growth inhibitor Cetermin Glioblastoma-derived T-cell suppressor factor G-TSF Polyergin Latency-associated peptide LAP
Function	TGF-beta 2 has suppressive effects on interleukin-2 dependent T-cell growth.
Cellular Localization	Secreted.
Post-translational Modifications	The precursor is cleaved into mature TGF-beta-2 and LAP, which remains non-covalently linked to mature TGF-beta-2 rendering it inactive.

St John's Laboratory Ltd

F +44 (0)207 681 2580

T +44 (0)208 223 3081

W <http://www.stjohnslabs.com/>

E info@stjohnslabs.com