


Anti-TNFAIP8L3 antibody



Description	Rabbit polyclonal to TNFAIP8L3.
Model	STJ98863
Host	Rabbit
Reactivity	Human, Mouse, Rat
Applications	ELISA, WB
Immunogen	Synthesized peptide derived from Human TNFAIP8L3
Immunogen Region	15-45 aa
Gene ID	388121
Gene Symbol	TNFAIP8L3
Dilution range	WB 1:500-2000ELISA 1:5000-20033
Specificity	This antibody detects endogenous levels of TNFAIP8L3.
Tissue Specificity	Widely expressed (at protein level) . Highly expressed in most carcinoma cell lines .
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Note	For Research Use Only (RUO).
Protein Name	Tumor necrosis factor alpha-induced protein 8-like protein 3 TNF alpha-induced protein 8-like protein 3 TNFAIP8-like protein 3
Clonality	Polyclonal
Conjugation	Unconjugated

Isotype	IgG
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Concentration	1 mg/ml
Storage Instruction	Store at -20°C, and avoid repeat freeze-thaw cycles.
Database Links	HGNC:20620OMIM:616438
Alternative Names	Tumor necrosis factor alpha-induced protein 8-like protein 3 TNF alpha-induced protein 8-like protein 3 TNFAIP8-like protein 3
Function	Acts as a lipid transfer protein. Preferentially captures and shuttles two lipid second messengers, i.e., phosphatidylinositol 4,5- bisphosphate and phosphatidylinositol 3,4,5-trisphosphate and increases their levels in the plasma membrane. Additionally, may also function as a lipid-presenting protein to enhance the activity of the PI3K-AKT and MEK-ERK pathways. May act as a regulator of tumorigenesis through its activation of phospholipid signaling.
Cellular Localization	Cytoplasm Cell membrane. On PDGF activation, translocates from cytoplasm to plasma membrane.

St John's Laboratory Ltd

F +44 (0)207 681 2580

T +44 (0)208 223 3081

W <http://www.stjohnslabs.com/>

E info@stjohnslabs.com