


Anti-CYCS antibody



Western Blot (WB) analysis of 1. HeLa 2. A549 cells using Anti-CYCS antibody. (STJ98953)



Description

CYCS is a protein encoded by the CYCS gene which is approximately 11,7 kDa. CYCS is localised to the mitochondrion intermembrane space. It is involved in the TNFR1 pathway, respiratory electron transport and apoptosis modulation and signalling. It is a small heme protein that functions as a central component of the electron transport chain in the mitochondria. It associates with the inner membrane of the mitochondrion where it accepts electrons from cytochrome b and transfers them to the cytochrome oxidase complex. It is also involved in the initiation of apoptosis. CYCS is expressed in the nervous system, liver, kidney, heart and muscle. Mutations in the CYCS gene may result in thrombocytopenia. STJ98953 was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. This primary antibody detects endogenous CYCS.

Model	STJ98953
Host	Rabbit
Reactivity	Human, Mouse, Rat
Applications	ELISA, WB
Immunogen	Synthetic peptide from human CYCS protein.
Immunogen Region	1-69 aa
Gene ID	54205
Gene Symbol	CYCS
Dilution range	WB 1:500-2000ELISA 1:10000-20000
Specificity	The antibody detects endogenous CYCS.

Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Note	For Research Use Only (RUO).
Protein Name	Cytochrome c
Molecular Weight	15kDa
Clonality	Polyclonal
Conjugation	Unconjugated
Isotype	IgG
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Concentration	1 mg/ml
Storage Instruction	Store at -20°C, and avoid repeat freeze-thaw cycles.
Database Links	HGNC:19986OMIM:123970
Alternative Names	Cytochrome c
Function	Electron carrier protein. The oxidized form of the cytochrome c heme group can accept an electron from the heme group of the cytochrome c1 subunit of cytochrome reductase. Cytochrome c then transfers this electron to the cytochrome oxidase complex, the final protein carrier in the mitochondrial electron-transport chain.; Plays a role in apoptosis. Suppression of the anti-apoptotic members or activation of the pro-apoptotic members of the Bcl-2 family leads to altered mitochondrial membrane permeability resulting in release of cytochrome c into the cytosol. Binding of cytochrome c to Apaf-1 triggers the activation of caspase-9, which then accelerates apoptosis by activating other caspases.
Cellular Localization	Mitochondrion intermembrane space. Loosely associated with the inner membrane.
Post-translational Modifications	Binds 1 heme group per subunit.; Phosphorylation at Tyr-49 and Tyr-98 both reduce by half the turnover in the reaction with cytochrome c oxidase, down-regulating mitochondrial respiration.