


Anti-Tubulin gamma antibody





Description Rabbit polyclonal to Tubulin gamma.

Model STJ98974

Host Rabbit

Reactivity Human, Mouse, Rat

Applications ELISA, WB

Immunogen Synthetic peptide from human Tubulin gamma protein.

Immunogen Region 380-440 aa

Gene ID <u>7283</u>

Gene Symbol <u>TUBG1</u>

Dilution range WB 1:500-2000ELISA 1:10000-20000

Specificity The antibody detects endogenous Tubulin gamma.

Purification The antibody was affinity-purified from rabbit antiserum by affinity-

chromatography using epitope-specific immunogen.

Note For Research Use Only (RUO).

Protein Name Tubulin gamma-1 chain Gamma-1-tubulin Gamma-tubulin complex

component 1 GCP-1

Clonality Polyclonal

Conjugation Unconjugated

Isotype IgG

Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.

Concentration 1 mg/ml

Store at -20°C, and avoid repeat freeze-thaw cycles. **Storage Instruction**

Database Links HGNC:12417OMIM:191135

Tubulin gamma-1 chain Gamma-1-tubulin Gamma-tubulin complex **Alternative Names**

component 1 GCP-1

Function Tubulin is the major constituent of microtubules. The gamma chain is found at

> microtubule organizing centers (MTOC) such as the spindle poles or the centrosome. Pericentriolar matrix component that regulates alpha/beta chain minus-end nucleation, centrosome duplication and spindle formation.

Cytoplasm, cytoskeleton, microtubule organizing center, centrosome **Cellular Localization**

Post-translational Phosphorylation at Ser-131 by BRSK1 regulates centrosome duplication, **Modifications**

possibly by mediating relocation of gamma-tubulin and its associated proteins

from the cytoplasm to the centrosome.

St John's Laboratory Ltd

F +44 (0)207 681 2580 T+44 (0)208 223 3081

W http://www.stjohnslabs.com/ E info@stjohnslabs.com