


Anti-MEK1 antibody



Description	Mouse monoclonal to MEK1.
Model	STJ99087
Host	Mouse
Reactivity	Human
Applications	ELISA, WB
Immunogen	Recombinant human MEK1 protein.
Gene ID	5605
Gene Symbol	MAP2K1
Dilution range	WB 1:500-2000ELISA 1:10000-20000
Specificity	This antibody detects endogenous levels of MEK1/2 and does not cross-react with related proteins.
Tissue Specificity	Widely expressed, with extremely low levels in brain.
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Clone ID	6F10-D7-G6
Note	For Research Use Only (RUO).
Protein Name	Dual specificity mitogen-activated protein kinase kinase 2 MAP kinase kinase 2 MAPKK 2 ERK activator kinase 2 MAPK/ERK kinase 2 MEK 2
Molecular Weight	45kDa
Clonality	Monoclonal

Conjugation	Unconjugated
Isotype	IgG1
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Concentration	1 mg/ml
Storage Instruction	Store at -20°C, and avoid repeat freeze-thaw cycles.
Database Links	HGNC:6842OMIM:601263
Alternative Names	Dual specificity mitogen-activated protein kinase kinase 2 MAP kinase kinase 2 MAPKK 2 ERK activator kinase 2 MAPK/ERK kinase 2 MEK 2
Function	Dual specificity protein kinase which acts as an essential component of the MAP kinase signal transduction pathway. Binding of extracellular ligands such as growth factors, cytokines and hormones to their cell-surface receptors activates RAS and this initiates RAF1 activation. RAF1 then further activates the dual-specificity protein kinases MAP2K1/MEK1 and MAP2K2/MEK2. Both MAP2K1/MEK1 and MAP2K2/MEK2 function specifically in the MAPK/ERK cascade, and catalyze the concomitant phosphorylation of a threonine and a tyrosine residue in a Thr-Glu-Tyr sequence located in the extracellular signal-regulated kinases MAPK3/ERK1 and MAPK1/ERK2, leading to their activation and further transduction of the signal within the MAPK/ERK cascade. Depending on the cellular context, this pathway mediates diverse biological functions such as cell growth, adhesion, survival and differentiation, predominantly through the regulation of transcription, metabolism and cytoskeletal rearrangements. One target of the MAPK/ERK cascade is peroxisome proliferator-activated receptor gamma (PPARG), a nuclear receptor that promotes differentiation and apoptosis. MAP2K1/MEK1 has been shown to export PPARG from the nucleus. The MAPK/ERK cascade is also involved in the regulation of endosomal dynamics, including lysosome processing and endosome cycling through the perinuclear recycling compartment (PNRC), as well as in the fragmentation of the Golgi apparatus during mitosis.
Sequence and Domain Family	The proline-rich region localized between residues 270 and 307 is important for binding to RAF1 and activation of MAP2K1/MEK1.
Cellular Localization	Cytoplasm, cytoskeleton, microtubule organizing center, centrosome Cytoplasm, cytoskeleton, microtubule organizing center, spindle pole body Cytoplasm Nucleus Membrane. Localizes at centrosomes during prometaphase, midzone during anaphase and midbody during telophase/cytokinesis . Membrane localization is probably regulated by its interaction with KSR1 .
Post-translational Modifications	Phosphorylation at Ser-218 and Ser-222 by MAP kinase kinase kinases (RAF or MEKK1) positively regulates kinase activity. Also phosphorylated at Thr-292 by MAPK1/ERK2 and at Ser-298 by PAK. MAPK1/ERK2 phosphorylation of Thr-292 occurs in response to cellular adhesion and leads to inhibition of Ser-298 phosphorylation by PAK. Acetylation by Yersinia yopJ prevents phosphorylation and activation, thus blocking the MAPK signaling pathway.

T +44 (0)208 223 3081

E info@stjohnslabs.com