


## Anti-Keratin 18 antibody



<b>Description</b>	Mouse monoclonal to Keratin 18.
<b>Model</b>	STJ99094
<b>Host</b>	Mouse
<b>Reactivity</b>	Human, Simian
<b>Applications</b>	ELISA, WB
<b>Immunogen</b>	Purified recombinant human Keratin 18 protein fragments expressed in E.coli.
<b>Gene ID</b>	<a href="#">3875</a>
<b>Gene Symbol</b>	<a href="#">KRT18</a>
<b>Dilution range</b>	WB 1:500-2000ELISA 1:10000-20000
<b>Specificity</b>	This antibody detects endogenous levels of Keratin 18 and does not cross-react with related proteins.
<b>Tissue Specificity</b>	Expressed in colon, placenta, liver and very weakly in exocervix. Increased expression observed in lymph nodes of breast carcinoma.
<b>Purification</b>	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
<b>Clone ID</b>	2B5-B6-F7
<b>Note</b>	For Research Use Only (RUO).
<b>Protein Name</b>	Keratin, type I cytoskeletal 18 Cell proliferation-inducing gene 46 protein Cytokeratin-18 CK-18 Keratin-18 K18
<b>Molecular Weight</b>	46kDa

<b>Clonality</b>	Monoclonal
<b>Conjugation</b>	Unconjugated
<b>Isotype</b>	IgG2a
<b>Formulation</b>	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
<b>Concentration</b>	1 mg/ml
<b>Storage Instruction</b>	Store at -20°C, and avoid repeat freeze-thaw cycles.
<b>Database Links</b>	<a href="#">HGNC:64300MIM:148070</a>
<b>Alternative Names</b>	Keratin, type I cytoskeletal 18 Cell proliferation-inducing gene 46 protein Cytokeratin-18 CK-18 Keratin-18 K18
<b>Function</b>	Involved in the uptake of thrombin-antithrombin complexes by hepatic cells . When phosphorylated, plays a role in filament reorganization. Involved in the delivery of mutated CFTR to the plasma membrane. Together with KRT8, is involved in interleukin-6 (IL-6)-mediated barrier protection.
<b>Cellular Localization</b>	Cytoplasm, perinuclear region. Nucleus, nucleolus.
<b>Post-translational Modifications</b>	Phosphorylation at Ser-34 increases during mitosis. Hyperphosphorylated at Ser-53 in diseased cirrhosis liver. Phosphorylation increases by IL-6. Proteolytically cleaved by caspases during epithelial cell apoptosis. Cleavage occurs at Asp-238 by either caspase-3, caspase-6 or caspase-7. O-GlcNAcylation increases solubility, and decreases stability by inducing proteasomal degradation.