


Anti-MMP-2 antibody



Description	Mouse monoclonal to MMP-2.
Model	STJ99213
Host	Mouse
Reactivity	Human, Rat
Applications	ELISA, WB
Immunogen	Purified recombinant human MMP-2 protein expressed in E.coli.
Gene ID	4313
Gene Symbol	MMP2
Dilution range	WB 1:500-2000ELISA 1:10000-20000
Specificity	This antibody detects endogenous levels of MMP-2 and does not cross-react with related proteins.
Tissue Specificity	Produced by normal skin fibroblasts. PEX is expressed in a number of tumors including gliomas, breast and prostate.
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Clone ID	3A8-C1-E8
Note	For Research Use Only (RUO).
Protein Name	72 kDa type IV collagenase 72 kDa gelatinase Gelatinase A Matrix metalloproteinase-2 MMP-2 TBE-1 PEX
Molecular Weight	72kDa

Clonality	Monoclonal
Conjugation	Unconjugated
Isotype	IgG1
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Concentration	1 mg/ml
Storage Instruction	Store at -20°C, and avoid repeat freeze-thaw cycles.
Database Links	HGNC:7166OMIM:120360
Alternative Names	72 kDa type IV collagenase 72 kDa gelatinase Gelatinase A Matrix metalloproteinase-2 MMP-2 TBE-1 PEX
Function	Ubiquitous metalloproteinase that is involved in diverse functions such as remodeling of the vasculature, angiogenesis, tissue repair, tumor invasion, inflammation, and atherosclerotic plaque rupture. As well as degrading extracellular matrix proteins, can also act on several nonmatrix proteins such as big endothelial 1 and beta-type CGRP promoting vasoconstriction. Also cleaves KISS at a Gly- -Leu bond. Appears to have a role in myocardial cell death pathways. Contributes to myocardial oxidative stress by regulating the activity of GSK3beta. Cleaves GSK3beta in vitro. Involved in the formation of the fibrovascular tissues in association with MMP14.; PEX, the C-terminal non-catalytic fragment of MMP2, possesses anti-angiogenic and anti-tumor properties and inhibits cell migration and cell adhesion to FGF2 and vitronectin. Ligand for integrin α v/ β 3 on the surface of blood vessels.; Isoform 2: Mediates the proteolysis of CHUK/IKKA and initiates a primary innate immune response by inducing mitochondrial-nuclear stress signaling with activation of the pro-inflammatory NF-kappaB, NFAT and IRF transcriptional pathways.
Sequence and Domain Family	The conserved cysteine present in the cysteine-switch motif binds the catalytic zinc ion, thus inhibiting the enzyme. The dissociation of the cysteine from the zinc ion upon the activation-peptide release activates the enzyme.
Cellular Localization	Isoform 1: Secreted, extracellular space, extracellular matrix. Membrane. Nucleus. Colocalizes with integrin α V/ β 3 at the membrane surface in angiogenic blood vessels and melanomas. Found in mitochondria, along microfibrils, and in nuclei of cardiomyocytes.. Isoform 2: Cytoplasm. Mitochondrion.
Post-translational Modifications	Phosphorylation on multiple sites modulates enzymatic activity. Phosphorylated by PKC in vitro. The propeptide is processed by MMP14 (MT-MMP1) and MMP16 (MT-MMP3). Autocatalytic cleavage in the C-terminal produces the anti-angiogenic peptide, PEX. This processing appears to be facilitated by binding integrin α v/ β 3.