


Anti-E2F-1 antibody



Description	Mouse monoclonal to E2F-1.
Model	STJ99250
Host	Mouse
Reactivity	Human, Rat
Applications	ELISA, WB
Immunogen	Purified recombinant human E2F-1 protein fragments expressed in E.coli.
Gene ID	1869
Gene Symbol	E2F1
Dilution range	WB 1:500-2000ELISA 1:10000-20000
Specificity	This antibody detects endogenous levels of E2F-1 and does not cross-react with related proteins.
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Clone ID	4G8-2C2-C12
Note	For Research Use Only (RUO).
Protein Name	Transcription factor E2F1 E2F-1 PBR3 Retinoblastoma-associated protein 1 RBAP-1 Retinoblastoma-binding protein 3 RBBP-3 pRB-binding protein E2F-1
Molecular Weight	70kDa
Clonality	Monoclonal

Conjugation	Unconjugated
Isotype	IgG2b
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Concentration	1 mg/ml
Storage Instruction	Store at -20°C, and avoid repeat freeze-thaw cycles.
Database Links	HGNC:31130MIM:189971
Alternative Names	Transcription factor E2F1 E2F-1 PBR3 Retinoblastoma-associated protein 1 RBAP-1 Retinoblastoma-binding protein 3 RBBP-3 pRB-binding protein E2F-1
Function	Transcription activator that binds DNA cooperatively with DP proteins through the E2 recognition site, 5'-TTTC[CG]CGC-3' found in the promoter region of a number of genes whose products are involved in cell cycle regulation or in DNA replication. The DRTF1/E2F complex functions in the control of cell-cycle progression from G1 to S phase. E2F1 binds preferentially RB1 in a cell-cycle dependent manner. It can mediate both cell proliferation and TP53/p53-dependent apoptosis. Blocks adipocyte differentiation by binding to specific promoters repressing CEBPA binding to its target gene promoters .
Cellular Localization	Nucleus.
Post-translational Modifications	Phosphorylated by CDK2 and cyclin A-CDK2 in the S-phase. Phosphorylation at Ser-364 by CHEK2 stabilizes E2F1 upon DNA damage and regulates its effect on transcription and apoptosis. Acetylation stimulates DNA-binding. Enhanced under stress conditions such as DNA damage and inhibited by retinoblastoma protein RB1. Regulated by KAP1/TRIM28 which recruits HDAC1 to E2F1 resulting in deacetylation. Acetylated by P/CAF/KAT2B.