

## Anti-Acetyl-Histone H4-K91 antibody (STJ11100937)

STJ11100937

### GENERAL INFORMATION


<b>Product Type</b>	Primary antibodies
<b>Short Description</b>	Rabbit polyclonal antibody anti-Acetyl-Histone H4-K91 is suitable for use in Western Blot and Immunofluorescence.
<b>Applications</b>	WB, IF
<b>Host/Source</b>	Rabbit
<b>Reactivity</b>	Human, Mouse, Rat

### PRODUCT PROPERTIES


<b>Clonality</b>	Polyclonal
<b>Clone ID</b>	
<b>Concentration</b>	
<b>Conjugation</b>	Unconjugated
<b>Purification</b>	Affinity purification
<b>Dilution Range</b>	WB 1:500-1:1000 IF 1:50-1:200
<b>Formulation</b>	PBS containing 0.02% Sodium Azide, 50% Glycerol, pH7.3.
<b>Isotype</b>	IgG
<b>Storage Instruction</b>	Store in a freezer at -20°C and avoid freeze-thaw cycles.

### TARGET INFORMATION


<b>Gene ID</b>	121504/554313/8294/8359/8360/8361/8362/8363/8364/8365/8366/8367/8368/8370
<b>Gene Symbol</b>	H4C1.H4C2.H4C3.H4C4.H4C5.H4C6.H4C8.H4C9.H4C11.H4C12.H4C13.H4C14.H4C15.H4-16
<b>Uniprot ID</b>	H4_HUMAN
<b>Immunogen</b>	A synthetic Peptide of human Acetyl-Histone H4-K91
<b>Immunogen Region</b>	
<b>Specificity</b>	
<b>Immunogen Sequence</b>	




Western blot analysis of extracts of NIH/3T3 cells, using Acetyl-Histone H4-K91 antibody (STJ11100937) at 1:1000 dilution. NIH/3T3 cells were treated by TSA (1  $\mu$ M) at 37 °C for 18 hours. Secondary antibody: HRP Goat Anti-rabbit IgG (H+L) at 1:10000 dilution. Lysates/proteins: 25 $\mu$ g per lane. Blocking buffer: 5% nonfat dry milk in TBST. Detection: ECL Basic Kit. Exposure time: 300s.



Immunofluorescence analysis of C6 cells using Acetyl-Histone H4-K91 antibody (STJ11100937) at dilution of 1:100. C6 cells were treated by TSA (1  $\mu$ M) at 37 °C for 18 hours. Blue: DAPI for nuclear staining.



Immunofluorescence analysis of HeLa cells using Acetyl-Histone H4-K91 antibody (STJ11100937) at dilution of 1:100. HeLa cells were treated by TSA (1  $\mu$ M) at 37 °C for 18 hours. Blue: DAPI for nuclear staining.



Immunofluorescence analysis of NIH/3T3 cells using Acetyl-Histone H4-K91 antibody (STJ11100937) at dilution of 1:100. NIH/3T3 cells were treated by TSA (1  $\mu$ M) at 37 °C for 18 hours. Blue: DAPI for nuclear staining.

This product is suitable for in-vitro studies under the RESEARCH USE ONLY [RUO] licence. This product must not be used as for diagnostic or other medical purposes.  
St John's Laboratory Ltd, Knowledge Dock Business Centre, University Way, London, E16 2RD | Tel: 0208 223 3081