

Anti-Acetyl-Histone H3-K56 antibody (STJ29398)

STJ29398

GENERAL INFORMATION


Product Type	Primary antibodies
Short Description	Rabbit polyclonal antibody anti-Acetyl-Histone H3-K56 is suitable for use in Western Blot, Immunohistochemistry, Immunofluorescence and Immunoprecipitation.
Applications	WB, IHC, IF, IP
Host/Source	Rabbit
Reactivity	Human, Mouse, Rat

PRODUCT PROPERTIES


Clonality	Polyclonal
Clone ID	
Concentration	
Conjugation	Unconjugated
Purification	Affinity purification
Dilution Range	WB 1:500-1:2000 IHC 1:50-1:200 IF 1:50-1:200 IP 1:50-1:200 ChIP 1:20-1:100 ChIP-seq 1:20-1:100
Formulation	PBS containing 0.02% Sodium Azide, 50% Glycerol, pH7.3.
Isotype	IgG
Storage	Store in a freezer at -20°C and avoid freeze-thaw cycles.
Instruction	

TARGET INFORMATION


Gene ID	8290
Gene Symbol	H3-4
Uniprot ID	H31T_HUMAN
Immunogen	A synthetic peptide of human Acetyl-Histone H3-K56
Immunogen Region	
Specificity	
Immunogen Sequence	




Western blot analysis of extracts of NIH/3T3 cells, using Acetyl-Histone H3-K56 antibody (STJ29398) at 1:1000 dilution. NIH/3T3 cells were treated by TSA (1 μ M) at 37 °C for 18 hours. Secondary antibody: HRP Goat Anti-rabbit IgG (H+L) at 1:10000 dilution. Lysates/proteins: 25ug per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit. Exposure time: 180s.



Immunofluorescence analysis of C6 cells using Acetyl-Histone H3-K56 rabbit polyclonal antibody (STJ29398) at dilution of 100 (40x lens). C6 cells were treated by TSA (1 μ M) at 37 °C for 18 hours. Blue: DAPI for nuclear staining.



Immunofluorescence analysis of NIH/3T3 cells using Acetyl-Histone H3-K56 rabbit polyclonal antibody (STJ29398) at dilution of 100 (40x lens). NIH/3T3 cells were treated by TSA (1 μ M) at 37 °C for 18 hours. Blue: DAPI for nuclear staining.



Immunofluorescence analysis of U-2 OS cells using Acetyl-Histone H3-K56 rabbit polyclonal antibody (STJ29398) at dilution of 100 (40x lens). U-2 OS cells were treated by TSA (1 μ M) at 37 °C for 18 hours. Blue: DAPI for nuclear staining.

This product is suitable for in-vitro studies under the RESEARCH USE ONLY [RUO] licence. This product must not be used as for diagnostic or other medical purposes.
St John's Laboratory Ltd, Knowledge Dock Business Centre, University Way, London, E16 2RD | Tel: 0208 223 3081