


MBM-55S

Chemical Properties

CAS No.: 2083624-07-9
Formula: C₃₆H₃₉FN₆O₁₀
Molecular Weight: 734.73
Appearance: N/A
Storage: 0-4°C for short term (days to weeks), or -20°C for long term (months).

**Biological Description**

Description	MBM-55S effectively inhibits the proliferation of cancer cells by inducing cell cycle arrest and apoptosis. MBM-55S shows antitumor activities, and no obvious toxicity to mice. MBM-55S is a potent NIMA-related kinase 2 (Nek2) inhibitor with an IC ₅₀ of 1 nM. MBM-55S shows a 20-fold or greater selectivity in most kinases with the exception of RSK1 (IC ₅₀ =5.4 nM) and DYRK1a (IC ₅₀ =6.5 nM).
Targets(IC ₅₀)	Nek2: 1 nM RSK1: 5.4 nM DYRK1a: 6.5 nM CHK1: 57 nM GSK-3β: 91 nM ABL: 20 nM CDK2: 370 nM CDK4: 441nM AKT1: 608 nM Aurora A: 5300 nM
In vitro	MBM-55S (0.5-1 μM; 24 hours) induces G2/M phase arrest and accumulation of cells with >4N content in HCT-116 cells. MBM-55S (0.5-1 μM; 24 hours) causes cell apoptosis in a concentration-dependent manner in HCT-116 cells. MBM-55S inhibits MGC-803, HCT-116, Bel-7402 cells proliferation with IC ₅₀ s of 0.53, 0.84, 7.13 μM, respectively.
In vivo	MBM-55S (1.0 mg/kg; i.v.) treatment shows the CL, V _{ss} , T _{1/2} , AUC _{0-t} , and AUC _{0-∞} values of 33.3 mL/min/kg, 2.53 L/kg, 1.72 hours, 495 ng/h/mL and 507 ng/h/mL, respectively. MBM-55S (20 mg/kg; i.p.; twice a day for 21 days) exhibits good antitumor activity and a well-tolerated dose schedule in nude mice bearing HCT-116 xenografts.

Solubility Information

Solubility	< 1 mg/ml refers to the product slightly soluble or insoluble
------------	---

Preparing Stock Solutions

	1mg	5mg	10mg
1 mM	1.361 mL	6.805 mL	13.61 mL
5 mM	0.272 mL	1.361 mL	2.722 mL
10 mM	0.136 mL	0.681 mL	1.361 mL
50 mM	0.027 mL	0.136 mL	0.272 mL

Please select the appropriate solvent to prepare the stock solution, according to the solubility of the product in different solvents. The storage conditions and period of the stock solution: - 80 °C for 6 months; - 20 °C for 1 month. Please use it as soon as possible.

Reference

1. Xi JB, et al. Structure-based design and synthesis of imidazo[1,2-a]pyridine derivatives as novel and potent Nek2 inhibitors with in vitro and in vivo antitumor activities. Eur J Med Chem. 2017 Jan 27;126:1083-1106.

[Inhibitors](#) · [Natural Compounds](#) · [Compound Libraries](#)

This product is for Research Use Only · Not for Human or Veterinary or Therapeutic Use.

Tel:781-999-4286

E-mail:info@targetmol.com

Address:36 Washington Street,Wellesley Hills,MA 02481