


Sulfacetamide sodium monohydrate

Chemical Properties

CAS No. :	6209-17-2
Formula:	C ₈ H ₉ N ₂ NaO ₃ S·H ₂ O
Molecular Weight:	254.2
Appearance:	no data available
Storage:	Powder: -20°C for 3 years In solvent: -80°C for 1 year



Biological Description

Description	Sulfacetamide sodium monohydrate, a sulfonamide antibiotic, has been investigated for the treatment of rosacea and pityriasis versicolor.
Targets(IC50)	Antibacterial,Antibiotic,Autophagy
In vitro	Sulfacetamide inhibits Arabidopsis DHPS with IC ₅₀ of 9.5 μM, pKa=5.4[1]. Sulfacetamide induces anti-proliferative effects on T-47D cells and it is independent of apoptosis and cell cycle arrest. Sulfacetamide treatment lowers expression of p53/DRAM pathway in parallel with upregulation of Akt/mTOR pathway promoting cytoprotective autophagy. The LD ₅₀ of sulfacetamide in T-47D cells after 48 h is 41 mM. Sulfacetamide does not cause DNA fragmentation. In cells treated with sulfacetamide, the ATG5 expression level increases suggesting an increase in autophagosome formation in the autophagy pathway. Autophagy induction in the sulfathiazole and sulfacetamide treatments is not accompanied by apoptosis and occurred without any distinctive arrest in a phase of the cell cycle. It triggers autophagy in T-47D cells via a DAPK independent pathway[2]. Sodium sulfacetamide or sulfacetamide is a bacteriostatic agent that is active against sulfonamide-sensitive Gramnegative and Gram-positive bacteria, including Streptococci, Staphylococci, E. coli, Klebsiella pneumoniae, Pseudomonas pyocyanea, Salmonella spp., Proteus vulgaris, and Nocardia, which are usually isolated in secondary infections of the skin. Sulfacetamide inhibits mannose-6-phosphate isomerase (also known as phosphomannose isomerase (PMI)), which is considered the key enzyme in kinetoplastid energy metabolism[3].
In vivo	The LD ₅₀ of sulfacetamide for mice is 16,500 mg/kg by the oral route. In humans, the side effects include erythema, moderate swelling, nausea, vomiting, and headache. In addition to these side effects, the occurrence of StevensJohnson syndrome is reported in HIV-positive patients who received sulfacetamide drops for eye infections. All of these side effects, however, are associated with oral administration or high drug absorption through the skin, mucous membranes, and the conjunctiva, whereas topical use is not associated with strong side effects.
Cell Research	Cells are cultured in RPMI medium 1640, supplemented with 10% FBS and 1% penicillin/streptomycin, in a humidified atmosphere of 5% carbon dioxide in air at 37°C. According to MTT assay, the LC ₅₀ of sulfathiazole and sulfacetamide after 48 h is determined as 6.5 mM and 41 mM, respectively. Doxorubicin and sodium salt of sulfadruugs are dissolved in culture medium to the final desired concentration based on the determined LC ₅₀ and filtered. Cells (at 80% confluency) are incubated with freshly

prepared drugs for 48h in a humidified incubator before being trypsinized and washed with phosphate-buffer saline 3 times and stored at 270°C. For cell viability assay, cells are seeded in at least triplicate wells for each concentration of drug per time at 1×10^4 cells/well in a 96-well plate. After 24h of seeding, the cells has grown to ~80% confluency. The medium is changed to that containing drugs at concentrations ranging from 0.0-50 mM. The concentration range for doxorubicin is 0-6 μ M. After 24, 48 and 72 h, each well is filled with 25 μ l MTT stock solution (4 mg/ml or 100 μ g/well) and incubated for 3 h at 37°C. Formazan crystals are dissolved in 100 μ l of dimethyl sulfoxide (DMSO) and quantified using a microplate reader at 570 nm. The MTT assays are performed at least 3 times for each drug and the percentage of surviving cells relative to control (untreated sample) is calculated.(Only for Reference)

Solubility Information

Solubility	DMSO: 16.67 mg/mL (65.57 mM),Sonication is recommended. H2O: 196.7 mM,Sonication is recommended. (< 1 mg/ml refers to the product slightly soluble or insoluble)
------------	--

Preparing Stock Solutions

	1mg	5mg	10mg
1 mM	3.9339 mL	19.6696 mL	39.3391 mL
5 mM	0.7868 mL	3.9339 mL	7.8678 mL
10 mM	0.3934 mL	1.967 mL	3.9339 mL
50 mM	0.0787 mL	0.3934 mL	0.7868 mL

Please select the appropriate solvent to prepare the stock solution, according to the solubility of the product in different solvents. Please use it as soon as possible.

Reference

Prabhu V, et al. Phytochemistry. 1997, 45(1):23-7.

Mohammadpour R, et al. Cell Biol Int. 2013, 37(4):348-58.

Lourival A. Silva, et al. BioMed Research International. 2015(2015), Article ID 965725.

Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins

This product is for Research Use Only· Not for Human or Veterinary or Therapeutic Use

Tel:781-999-4286 E_mail:info@targetmol.com Address:36 Washington Street,Wellesley Hills,MA 02481