


Diazoxide

Chemical Properties

CAS No. :	364-98-7
Formula:	C ₈ H ₇ ClN ₂ O ₂ S
Molecular Weight:	230.67
Appearance:	no data available
Storage:	Powder: -20°C for 3 years In solvent: -80°C for 1 year



Biological Description

Description	Diazoxide (Proglycem) is a benzothiadiazine derivative that is a peripheral vasodilator used for hypertensive emergencies. It lacks diuretic effect, apparently because it lacks a sulfonamide group.
Targets(IC50)	ATPase, Autophagy, Potassium Channel
In vitro	Diazoxide inhibits microglial inflammatory activity. Diazoxide treatment partially inhibits the inflammatory pattern induced by LPS/IFN- γ in microglial cells, inducing a decrease in NO production that could be because of the decreased expression of iNOS detected. Diazoxide has no effect on microglial phagocytosis[1].
In vivo	Diazoxide is beneficial on the improvement in cognitive tasks, reduction of anxiety, decrease in the accumulation of amyloid-beta oligomers and hyperphosphorylation of tau proteins. Diazoxide may also exerts neuroprotective effects independently of K ⁺ channel activation by decreasing neuronal excitability and activation of N-methyl-D-aspartate (NMDA) receptors or by increasing currents through α -amino-3-hydroxy-5-methyl-4-isoxazolepropionic acid (AMPA) receptors. Diazoxide-treated animals show a decrease in disease severity a few days after the first clinical signs are observed, corresponding to the acute inflammatory phase of the disease. Daily oral administration of diazoxide in EAE mice during the effector phase of the disease reduces the severity of the clinical signs without any apparent adverse effect. Diazoxide decreases demyelination and axonal loss, reduces tissue damage, inhibits microglial/macrophage and astrocytic activation and preserves neuron integrity. No effects are observed on the number of B and T lymphocytes infiltrating the spinal cord[1].
Cell Research	The phagocytic ability of microglia is determined by the uptake of 2- μ m red fluorescent microspheres by BV-2 cells. Cells are treated with diazoxide 100 μ M and activated with LPS/IFN- γ and then incubated with microspheres at a concentration of 0.01% for 30 min in the dark at 37°C and 5% CO ₂ . Cells are rinsed twice in PBS solution, pelleted at 1,000 g for 5 min and resuspended in 300 μ L PBS. Cells are kept on ice and analyzed by flow cytometry.(Only for Reference)

Solubility Information

Solubility	H ₂ O: < 1 mg/mL (insoluble or slightly soluble), Ethanol: < 1 mg/mL (insoluble or slightly soluble), DMSO: 60 mg/mL (260.11 mM), Sonication is recommended.
------------	---

(< 1 mg/ml refers to the product slightly soluble or insoluble)

Preparing Stock Solutions

	1mg	5mg	10mg
1 mM	4.3352 mL	21.676 mL	43.352 mL
5 mM	0.867 mL	4.3352 mL	8.6704 mL
10 mM	0.4335 mL	2.1676 mL	4.3352 mL
50 mM	0.0867 mL	0.4335 mL	0.867 mL

Please select the appropriate solvent to prepare the stock solution, according to the solubility of the product in different solvents. Please use it as soon as possible.

Reference

Virgili N, et al. J Neuroinflammation. 2011, 8:149.
Garlid KD, et al. Circ Res. 1997, 81(6):1072-1082.

Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins

This product is for Research Use Only· Not for Human or Veterinary or Therapeutic Use

Tel:781-999-4286 E_mail:info@targetmol.com Address:36 Washington Street,Wellesley Hills,MA 02481