


Vanillylmandelic acid

Chemical Properties

CAS No. :	55-10-7
Formula:	C ₉ H ₁₀ O ₅
Molecular Weight:	198.17
Appearance:	no data available
Storage:	Powder: -20°C for 3 years In solvent: -80°C for 1 year



Biological Description

Description	Vanillylmandelic acid, the end product of the metabolism of catecholamines, epinephrine, and norepinephrine, has antioxidant activity and is commonly used to help diagnose pheochromocytoma, as well as its elevation in association with depressive symptoms.
Targets(IC50)	Antioxidant
Kinase Assay	In Vitro Protein-tyrosine Kinase Assays: Recombinant Syk is expressed in baculovirus-infected St9 cells. Assays of recombinant Syk activity are carried out using angiotensin I peptide as substrate. The enzyme activities of recombinant Syk are measured by phosphorylation of angiotensin I peptide in the presence of various concentrations of Piceatannol.

Solubility Information

Solubility	DMSO: 45 mg/mL (227.08 mM), Sonication is recommended. (< 1 mg/ml refers to the product slightly soluble or insoluble)
------------	---

Preparing Stock Solutions

	1mg	5mg	10mg
1 mM	5.0462 mL	25.2309 mL	50.4617 mL
5 mM	1.0092 mL	5.0462 mL	10.0923 mL
10 mM	0.5046 mL	2.5231 mL	5.0462 mL
50 mM	0.1009 mL	0.5046 mL	1.0092 mL

Please select the appropriate solvent to prepare the stock solution, according to the solubility of the product in different solvents. Please use it as soon as possible.

Reference

Kolentinis MK, et al. Eur J Pharmacol. 2013 Mar 5;703(1-3):46-52.

Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins

This product is for Research Use Only· Not for Human or Veterinary or Therapeutic Use

Tel:781-999-4286 E_mail:info@targetmol.com Address:36 Washington Street,Wellesley Hills,MA 02481