


## Nicotinamide N-oxide

## Chemical Properties

CAS No. :	1986-81-8
Formula:	C <sub>6</sub> H <sub>6</sub> N <sub>2</sub> O <sub>2</sub>
Molecular Weight:	138.12
Appearance:	no data available
Storage:	Powder: -20°C for 3 years   In solvent: -80°C for 1 year



## Biological Description

Description	Nicotinamide N-oxide (Nicotinamide 1-oxide) is recognized as an in vivo metabolite of nicotinamide which is a precursor of nicotinamide-adenine dinucleotide (NAD <sup>+</sup> ) in animals. The enzyme that catalyzes the reduction of nicotinamide N-oxide to nicotinamide in the liver is xanthine oxidase.
Targets(IC <sub>50</sub> )	Endogenous Metabolite,c-Myc,CXCR,Drug Metabolite
In vitro	Nicotinamide, a vitamin B3 variant, serves as a precursor to nicotinamide adenine dinucleotide, crucial for oxidative phosphorylation and dehydrogenase cofactor roles. It undergoes metabolism via two pathways: initial methylation by nicotinamide N-methyltransferase, followed by aldehyde oxidase oxidation, and direct oxidation to nicotinamide N-oxide. Studies have introduced a range of nicotinamide N-oxides as innovative, effective CXCR2 receptor antagonists. Specifically, Compound 1 effectively inhibits neutrophil chemotaxis (IC <sub>50</sub> =10 nM), whereas Compound 2 selectively blocks IL-8 binding (IC <sub>50</sub> =110 nM) and also significantly hampers neutrophil chemotaxis (IC <sub>50</sub> =170 nM).

## Solubility Information

Solubility	DMSO: 3.85 mg/mL (27.85 mM),Sonication is recommended. H <sub>2</sub> O: Limited solubility, (< 1 mg/ml refers to the product slightly soluble or insoluble)
------------	--

## Preparing Stock Solutions

	1mg	5mg	10mg
1 mM	7.2401 mL	36.2004 mL	72.4008 mL
5 mM	1.448 mL	7.2401 mL	14.4802 mL
10 mM	0.724 mL	3.620 mL	7.2401 mL
50 mM	0.1448 mL	0.724 mL	1.448 mL

Please select the appropriate solvent to prepare the stock solution, according to the solubility of the product in different solvents. Please use it as soon as possible.

### Reference

Cutshall NS, et al. Nicotinamide N-oxides as CXCR2 antagonists. Bioorg Med Chem Lett. 2001 Jul 23;11(14):1951-4.  
Real AM, et al. Nicotinamide N-oxidation by CYP2E1 in human liver microsomes. Drug Metab Dispos. 2013 Mar;41(3):550-3.

**Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins**

This product is for Research Use Only· Not for Human or Veterinary or Therapeutic Use

Tel:781-999-4286    E\_mail:info@targetmol.com    Address:36 Washington Street,Wellesley Hills,MA 02481