Data Sheet (Cat.No.T1048)



Estradiol

Chemical Properties

CAS No.: 50-28-2

Formula: C18H24O2

Molecular Weight: 272.38

Appearance: no data available

Storage: keep away from direct sunlight

torage: Powder: -20°C for 3 years | In solvent: -80°C for 1 year

он 0 _{На}С

Biological Description

Description

	fertility and maintenance of secondary sexual characteristics. Estradiol upregulates IL-6 expression through estrogen receptor beta (ERβ).				
Targets(IC50)	Estrogen Receptor/ERR,Estrogen/progestogen Receptor,Endogenous Metabolite				
In vitro	METHODS: Human lens epithelial cells HLEC were treated with Estradiol (0.001-10 μM) for 24 h. Cell viability was measured using the MTT. RESULTS: The number of viable cells was significantly reduced in the presence of 10 μM Estradiol. [1] METHODS: Human breast cancer cells MB231 were treated with Estradiol (100 nM) for 24 h. Cell motility was detected using scratch/wound healing assay and transwell assay. RESULTS: Estradiol treatment reduced the migration and invasion of MB231 cells. [2]				
In vivo	METHODS: To assay in vivo activity, Estradiol (0.2 mg/kg) and progesterone (5-20 mg/kg) were administered intraperitoneally as a single injection to ovariectomized C57BL/6 female mice. RESULTS: Mice receiving Estradiol alone or Estradiol plus 10-20 mg/kg progesterone showed significantly enhanced memory for new objects. [3] METHODS: To assay in vivo activity, Estradiol (0.2 mg/kg) was administered by gavage to ovariectomized C57BL/6 female mice once daily for two weeks. RESULTS: Long-term activation of ERβ reduced hot flash-like symptoms and enhanced spatial and object recognition memory in ovariectomized mice. [4]				
Cell Research	Estradiol is dissolved in 0.01% ethanol. Briefly, B9 cells (5×103 /well of a 96-well plate) are cultured with a series of dilutions of the supernatants in a final volume of 100 mL of RPMI 1640, supplemented with 5×10 -5mol/Lof 2-mercaptoethanol, 10% FBS, 100 U/mL penicillin, and 100 µg/mL streptomycin in flat-bottom microtiter plates. After 42 h, 0.5 µCi of [3H]thymidine is added. 6 h later, the cells are harvested and the radioactivity incorporated is determined. IL-6 is quantitated from a standard curve set up with known amounts of recombinant human or mouse IL-6. The anti-IL-6 monoclonal antibody completely inhibited the ability of the B9 cells to proliferate in response to recombinant mouse IL-6. In addition, we determined that B9 cells do not proliferate in response to IL-1 or TNF, and that antibodies to these two cytokines did not affect the cell proliferation in response to IL-6. Further, it is established that neither Estradiol (17β -estradiol), nor the 0.01% ethanol used as vehicle in these experiments, has any effect on the bioassay.				

Estradiol (E2) is a naturally occurring steroidal sex hormone that is essential for female

Page 1 of 3 www.targetmol.com

Solubility Information

Solubility	Ethanol: 20.4 mg/mL (74.9 mM), Sonication is recommended.
	DMSO: 50 mg/mL (183.57 mM), Sonication is recommended.
	10% DMSO+40% PEG300+5% Tween 80+45% Saline: 2.72 mg/mL (9.99 mM),Suspension.
	(< 1 mg/ml refers to the product slightly soluble or insoluble)

Preparing Stock Solutions

9-0	1mg	5mg	10mg	
1 mM	3.6713 mL	18.3567 mL	36.7134 mL	
5 mM	0.7343 mL	3.6713 mL	7.3427 mL	
10 mM	0.3671 mL	1.8357 mL	3.6713 mL	
50 mM	0.0734 mL	0.3671 mL	0.7343 mL	

Please select the appropriate solvent to prepare the stock solution, according to the solubility of the product in different solvents. Please use it as soon as possible.

Page 2 of 3 www.targetmol.com

Reference

Celojevic D, et al. Effects of 17β -estradiol on proliferation, cell viability and intracellular redox status in native human lens epithelial cells. Mol Vis. 2011;17:1987-96. Epub 2011 Jul 20.

Lu Z, Cao Y, Zhang D, et al. Discovery of Thieno [2, 3-e] indazole Derivatives as Novel Oral Selective Estrogen Receptor Degraders with Highly Improved Antitumor Effect and Favorable Druggability. Journal of Medicinal Chemistry. 2022

Xiao Y, Tao M, He J, et al.TIG1 Triggers Placental Senescence in Preeclampsia through LMNA/p53 Axis Activation. Placenta.2024

Liang L, Shen Z, Zhang K, et al.DLGAP5 upregulates E2F1 to promote prostate adenocarcinoma neuroendocrine differentiation.Life Sciences.2025: 123442.

Zhao Z, et al. 17β -Estradiol treatment inhibits breast cell proliferation, migration and invasion by decreasing MALAT-1 RNA level. Biochem Biophys Res Commun. 2014 Mar 7;445(2):388-93.

Qiao L, Wang H, He J, et al. Truncated affinity-improved aptamers for 17β-estradiol determination by AuNPs-based colorimetric aptasensor. Food Chemistry. 2020: 128181

Harburger LL, et al. Dose-dependent effects of post-training estradiol plus progesterone treatment on object memory consolidation and hippocampal extracellular signal-regulated kinase activation in young ovariectomized mice. Neuroscience. 2009 Apr 21;160(1):6-12.

Zhang S, Sun C, Zhao S, et al. Exposure to DEHP or its metabolite MEHP promotes progesterone secretion and inhibits proliferation in mouse placenta or JEG-3 cells. Environmental Pollution. 2020, 257: 113593
Fleischer AW, et al. Long-term oral administration of a novel estrogen receptor beta agonist enhances memory and alleviates drug-induced vasodilation in young ovariectomized mice. Horm Behav. 2021 Apr;130:104948.
Liu A, Jin M, Xie L, et al. Loss of miR-29a impairs decidualization of endometrial stromal cells by TET3 mediated demethylation of Col1A1 promoter. Iscience. 2021, 24(9): 103065.

Woolley CS, et al. J Comp Neurol, 1993, 336(2), 293-306.

Wang Y, Xu G, Wang Z, et al. Psoralidin, a major component of Psoraleae Fructus, induces inflammasome activation and idiosyncratic liver injury. International Immunopharmacology. 2021 Mar;92:107352. doi: 10.1016/j. intimp.2020.107352. Epub 2021 Jan 7.

Cui L, Xu F, Jiang Z, et al. Melatonin regulates proliferation and apoptosis of endometrial stromal cells via MT1. Acta Biochimica et Biophysica Sinica. 2021

Wang Y, Xu G, Wang Z, et al. Psoralidin, a major component of Psoraleae Fructus, induces inflammasome activation and idiosyncratic liver injury[J]. International Immunopharmacology. 2021, 92: 107352.

Zhang S, Sun C, Zhao S, et al. Exposure to DEHP or its metabolite MEHP promotes progesterone secretion and inhibits proliferation in mouse placenta or JEG-3 cells[J]. Environmental Pollution. 2020, 257: 113593.

Cui L, Xu F, Xu C, et al. Circadian gene Rev-erb α influenced by sleep conduces to pregnancy by promoting endometrial decidualization via IL-6-PR-C/EBP β axis. Journal of Biomedical Science. 2022, 29(1): 1-14.

Zhang D, Ma B, Liu D, et al.Discovery of a peptide proteolysis-targeting chimera (PROTAC) drug of p300 for prostate cancer therapy. EBio Medicine. 2024, 105.

Qiao L, Wang H, He J, et al. Truncated affinity-improved aptamers for 17β-estradiol determination by AuNPs-based colorimetric aptasensor[J]. Food Chemistry. 2020: 128181.

Zhang D, Lu Z, He Y, et al.Discovery of Thiochroman Derivatives as Potent, Oral Selective Estrogen Receptor Degraders and Antagonists for the Treatment of Endocrine-Resistant Breast Cancer. Journal of Medicinal Chemistry. 2024

Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins

This product is for Research Use Only. Not for Human or Veterinary or Therapeutic Use

Tel:781-999-4286 E_mail:info@targetmol.com Address:36 Washington Street,Wellesley Hills,MA 02481

Page 3 of 3 www.targetmol.com