

## Deramciclanc

## Chemical Properties


CAS No. : 120444-71-5

Formula: C<sub>20</sub>H<sub>31</sub>NO

Molecular Weight: 301.47

Appearance:

Storage: Powder: -20°C for 3 years | In solvent: -80°C for 1 year



## Biological Description

Description	Deramciclanc has high affinity for 5-HT <sub>2A</sub> and 5-HT <sub>2C</sub> receptors. It acts as an antagonist in both receptor subtypes and has inverse agonist properties for 5-HT <sub>2C</sub> receptors, but no direct stimulant agonists.
Targets(IC <sub>50</sub> )	5-HT Receptor
In vitro	Deramciclanc is a new anti-anxiety drug that can bind to 5-HT <sub>2A/2C</sub> receptor with high affinity. The interaction between deracixan and serotonin 5-HT <sub>2C</sub> receptor was further characterized using receptor phosphoinositide hydrolysis assay and receptor autoradiography. Deramciclanc antagonizes 5-HT <sub>2C</sub> receptor-mediated 5-HT stimulated phosphoinositide hydrolysis with IC <sub>50</sub> value of 168 nM. In the physiological system of the choroid plexus nervous system, deraciclanc can also reduce the hydrolysis of basic inositol phosphate by as much as 33% (EC <sub>50</sub> = 93 nM), which indicates that deraciclanc has inverse agonism at this receptor Agent properties.
In vivo	Compared with basal levels, at any time point, deramamine 3 and 10 mg/kg did not significantly change dopamine levels, while at 40-100 minutes and 160-240 minutes, delamethasone 30 mg/kg was significant Increased dopamine levels (P <0.05). Deramciclanc is a putative anti-serotonergic compound that reduces 5-HT-induced phosphoinositide hydrolysis and multiple effects caused by serotonergic agonists. The receptor binding curve of Deramciclanc is comparable to that of ritanserin. Deramciclanc has high affinity for 5-HT <sub>2A</sub> and 5-HT <sub>2C</sub> receptors. It acts as an antagonist in both receptor subtypes and has inverse agonist properties for 5-HT <sub>2C</sub> receptors without direct stimulatory agonist effects. In some animal experiments, deraciclanc has been shown to have anxiolytic effects.

### Preparing Stock Solutions

	1mg	5mg	10mg
1 mM	3.3171 mL	16.5854 mL	33.1708 mL
5 mM	0.6634 mL	3.3171 mL	6.6342 mL
10 mM	0.3317 mL	1.6585 mL	3.3171 mL
50 mM	0.0663 mL	0.3317 mL	0.6634 mL

Please select the appropriate solvent to prepare the stock solution, according to the solubility of the product in different solvents. Please use it as soon as possible.

### Reference

P?lvim?ki EP, et al. Deramciclane, a putative anxiolytic drug, is a serotonin 5-HT<sub>2C</sub> receptor inverse agonist but fails to induce 5-HT<sub>2C</sub> receptor down-regulation. *Psychopharmacology (Berl)*. 1998 Mar;136(2):99-104.

K??ri?inen TM, et al. Comparison of the effects of deramciclane, ritanserin and buspirone on extracellular dopamine and its metabolites in striatum and nucleus accumbens of freely moving rats. *Basic Clin Pharmacol Toxicol*. 2008 Jan;102(1):50-8.

**Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins**

**This product is for Research Use Only· Not for Human or Veterinary or Therapeutic Use**

Tel:781-999-4286    E\_mail:info@targetmol.com    Address:34 Washington Street,Wellesley Hills,MA 02481