


Dihydroergocristine mesylate

Chemical Properties

CAS No. :	24730-10-7
Formula:	C36H45N5O8S
Molecular Weight:	707.84
Appearance:	no data available
Storage:	Powder: -20°C for 3 years In solvent: -80°C for 1 year



Biological Description

Description	Dihydroergocristine mesylate (DHEC (mesylate))(DHEC mesylate) is the methanesulfonic acid salt of dihydroergocristine. It has been used as the for the symptomatic treatment of mental deterioration associated with cerebrovascular insufficiency and in peripheral vascular disease. Dihydroergocristine mesylate is a inhibitor of γ -secretase (GSI), , reduces the production of the Alzheimer's disease amyloid- β peptides, binds directly to γ -secretase and Nicastrin with equilibrium dissociation constants (Kd) of 25.7 nM and 9.8 μ M, respectively[1]. It is also a component of ergoloid mesylate (codergocrine mesilate), a mixture of ergot alkaloid derivatives that is used as a vasodilator and has shown mild benefits in the treatment of vascular dementia. It has a role as a vasodilator agent, an alpha-adrenergic antagonist and a geroprotector. It contains a dihydroergocristine.
Targets(IC50)	Beta Amyloid
In vitro	The IC50 value(2-20 μ M; 24 hours) of Dihydroergocristine (DHEC) for inhibiting the activity of γ -secretase in T100 cells without affecting cell viability is 25 μ M. Dihydroergocristine (2-20 μ M; 24 hours) inhibits cellular A β production and causes a dose-dependent accumulation of carboxy-terminal fragments of APP (APP-CTFs) in HEK293 and decreases γ -secretase activity in fibroblast cells[1].
Cell Research	Dihydroergocristine (2-20 μ M; 24 hours) causes a dose-dependent accumulation of carboxy-terminal fragments of APP (APP-CTFs) in HEK293 and decreases γ -secretase activity in fibroblast cells and inhibits cellular A β production[1].

Solubility Information

Solubility	DMSO: 60 mg/mL (84.76 mM),Sonication is recommended. H2O: 1 mg/mL (1.41 mM),Sonication is recommended. (< 1 mg/ml refers to the product slightly soluble or insoluble)
------------	--

Preparing Stock Solutions

	1mg	5mg	10mg
1 mM	1.4127 mL	7.0637 mL	14.1275 mL
5 mM	0.2825 mL	1.4127 mL	2.8255 mL
10 mM	0.1413 mL	0.7064 mL	1.4127 mL
50 mM	0.0283 mL	0.1413 mL	0.2825 mL

Please select the appropriate solvent to prepare the stock solution, according to the solubility of the product in different solvents. Please use it as soon as possible.

Reference

Lei X, et al. The FDA-approved natural product dihydroergocristine reduces the production of the Alzheimer's disease amyloid- β peptides. Sci Rep. 2015 Nov 16;5:16541.

Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins

This product is for Research Use Only. Not for Human or Veterinary or Therapeutic Use

Tel: 781-999-4286 E_mail: info@targetmol.com Address: 36 Washington Street, Wellesley Hills, MA 02481