


EN6

Chemical Properties

CAS No. : 1808714-73-9
 Formula: C₁₉H₁₄F₂N₄O₂
 Molecular Weight: 368.34
 Appearance: no data available
 Storage: Powder: -20°C for 3 years | In solvent: -80°C for 1 year



Biological Description

Description	EN6-mediated ATP6V1A modification decouples the v-ATPase from the Rags, leading to inhibition of mTORC1 signaling, increased lysosomal acidification and activation of autophagy. EN6 clears TDP-43 aggregates, a causative agent in frontotemporal dementia, in a lysosome-dependent manner. EN6 is a small-molecule in vivo activator of autophagy that covalently targets cysteine 277 in the ATP6V1A subunit of the lysosomal the vacuolar H ⁺ ATPase (v-ATPase).
Targets(IC50)	Proton pump,Autophagy

Solubility Information

Solubility	DMSO: 4.63 mg/mL (12.57 mM),Sonication is recommended. Ethanol: 1.10 mg/mL (2.99 mM),Sonication and heating to 60°C are recommended. (< 1 mg/ml refers to the product slightly soluble or insoluble)
------------	--

Preparing Stock Solutions

	1mg	5mg	10mg
1 mM	2.7149 mL	13.5744 mL	27.1488 mL
5 mM	0.543 mL	2.7149 mL	5.4298 mL
10 mM	0.2715 mL	1.3574 mL	2.7149 mL
50 mM	0.0543 mL	0.2715 mL	0.543 mL

Please select the appropriate solvent to prepare the stock solution, according to the solubility of the product in different solvents. Please use it as soon as possible.

Reference

Chung CY, et al. Covalent targeting of the vacuolar H⁺-ATPase activates autophagy via mTORC1 inhibition. Nat Chem Biol. 2019 Aug;15(8):776-785.

Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins

This product is for Research Use Only· Not for Human or Veterinary or Therapeutic Use

Tel:781-999-4286 E_mail:info@targetmol.com Address:36 Washington Street,Wellesley Hills,MA 02481