

FIDAS-5

Chemical Properties


CAS No. : 1391934-98-7

Formula: C₁₅H₁₃ClFN

Molecular Weight: 261.72

Appearance: no data available

Storage: Powder: -20°C for 3 years | In solvent: -80°C for 1 year



Biological Description

Description	FIDAS-5 is an orally active methionine adenosyltransferase 2A (MAT2A) inhibitor with an IC ₅₀ of 2.1 μM. It can effectively compete with S-adenosylmethionine for MAT2A binding and has anticancer effects.
Targets(IC ₅₀)	Others
In vitro	<p>METHODS: FIDAS-5 (3 μM; 36 hours) was used to treat LS174T cells, and the levels of S-adenosylmethionine (SAM) and S-adenosylhomocysteine (SAH) were observed.</p> <p>RESULTS FIDAS-5 reduced S-adenosylmethionine (SAM) and S-adenosylhomocysteine (SAH) levels in LS174T cells.</p> <p>METHODS: FIDAS-5 (3 μM; 7 days) was used to treat LS174T cells, and the proliferation of LS174T cells was observed.</p> <p>RESULTS FIDAS-5 significantly inhibited the proliferation of LS174T cells.</p> <p>METHODS: FIDAS-5 (3 μM) treatment inhibited LS174T colorectal cancer cells, and the expression of c-Myc and cyclinD1 in LS174T colorectal cancer cells was observed by westernBlot.</p> <p>RESULTS FIDAS-5 inhibited the expression of c-Myc and cyclinD1 in LS174T colorectal cancer cells [1].</p> <p>METHODS: OPM2 cells were transduced with siMAT2A or scrambled siRNA for 3 days. MAT2A-silenced and parental OPM2 cells were treated with different doses of FIDAS-5 (0.5, 1, 2 μM) to determine whether FIDAS-5 indeed impairs MM cell survival by inhibiting the enzymatic activity of MAT2A.</p> <p>RESULTS The antitumor effect of FIDAS-5 on MAT2A-silenced cells was impaired compared with cells transduced with scrambled siRNA. FIDAS-5 indeed inhibited MM cell survival in part by targeting MAT2A. [2]</p>
In vivo	<p>METHODS: Tumors were induced in athymic nude mice by subcutaneous injection of HT29 CRC cells. FIDAS-5 treatment (20 mg/kg) was administered orally via gavage. The oral efficacy of FIDAS-5 on HT29 tumor xenografts in nude mice was tested by measuring tumors twice weekly using digital calipers.</p> <p>RESULTS FIDAS-5 significantly inhibited xenograft tumor growth with minimal body weight differences. [1]</p>

Solubility Information

A DRUG SCREENING EXPERT

Solubility	DMSO: 125 mg/mL (477.61 mM),Sonication is recommended. (< 1 mg/ml refers to the product slightly soluble or insoluble)
------------	---

Preparing Stock Solutions

	1mg	5mg	10mg
1 mM	3.8209 mL	19.1044 mL	38.2088 mL
5 mM	0.7642 mL	3.8209 mL	7.6418 mL
10 mM	0.3821 mL	1.9104 mL	3.8209 mL
50 mM	0.0764 mL	0.3821 mL	0.7642 mL

Please select the appropriate solvent to prepare the stock solution, according to the solubility of the product in different solvents. Please use it as soon as possible.

Reference

Sundberg TB,et al. Development of Chemical Probes for Investigation of Salt-Inducible Kinase Function in Vivo. ACS Chem Biol. 2016 Aug 19;11(8):2105-11.

He X C, Wang J, Shi M Y, et al.Hypoxia-induced one-carbon metabolic reprogramming in glioma stem-like cells.Life Medicine.2023: lnad048.

Wang Y, et al. S-adenosylmethionine biosynthesis is a targetable metabolic vulnerability in multiple myeloma. Haematologica. 2024 Jan 1;109(1):256-271.

Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins

This product is for Research Use Only· Not for Human or Veterinary or Therapeutic Use

Tel:781-999-4286 E_mail:info@targetmol.com Address:36 Washington Street,Wellesley Hills,MA 02481