Data Sheet (Cat.No.T11684)



Isoguvacine hydrochloride

Chemical Properties


CAS No.: 68547-97-7

Formula: C6H10ClNO2

Molecular Weight: 163.6

Appearance: no data available

Storage: Powder: -20°C for 3 years | In solvent: -80°C for 1 year



HCI

Biological Description

Description	Isoguvacine hydrochloride (1,2,3,6-tetrahydro-4-pyridinecarboxylic acid hydrochloride) is a GABA receptor agonist. Isoguvacine hydrochloride binds to rat synaptic cortical membranes and activates $\alpha 1\beta 2\gamma 2S$, $\alpha 2\beta 2\gamma 2S$, $\alpha 3\beta 2\gamma 2S$, $\alpha 5\beta 2\gamma 2S$, and $\alpha 1S$ subunit GABAA receptors.
Targets(IC50)	GABA Receptor
In vitro	Treatment of 50 µM Isoguvacine to block epileptiform events in 2 of 6 organotypic hippocampal slice cultures, Isoguvacine dose-dependently inhibits hypomagnesium-induced epileptiform events[2]. Isoguvacine binds to mouse forebrain synaptic membrane preparations and can specifically bind to GABA, muscimol and bicuculline, but not picrotoxin or diaminobutyric acid. The highest levels of binding are observed in the cerebellum, cortex and hippocampus[3].

Solubility Information

Solubility	H2O: 90 mg/mL (550.12 mM), Sonication is recommended.	H2O: 90 mg/mL (550.12 mM),Sonication is recommended.	
	DMSO: 20 mg/mL (122.25 mM), Sonication is recommended.		
	(< 1 mg/ml refers to the product slightly soluble or insoluble)		

Preparing Stock Solutions

	1mg	5mg	10mg
1 mM	6.1125 mL	30.5623 mL	61.1247 mL
5 mM	1.2225 mL	6.1125 mL	12.2249 mL
10 mM	0.6112 mL	3.0562 mL	6.1125 mL
50 mM	0.1222 mL	0.6112 mL	1.2225 mL

Please select the appropriate solvent to prepare the stock solution, according to the solubility of the product in different solvents. Please use it as soon as possible.

Page 1 of 2 www.targetmol.com

Reference

White WF, et al. Isoguvacine binding, uptake, and release: relation to the GABA system. J Neurochem. 1983 Jun;40 (6):1701-8.

Wahab A, et al. Effects of gamma-aminobutyric acid (GABA) agonists and a GABA uptake inhibitor on pharmacoresistant seizure like events in organotypic hippocampal slice cultures. Epilepsy Res. 2009 Oct;86(2-3): 113-23.

Morin AM, et al. The binding of 3H-isoguvacine to mouse brain synapticmembranes. Life Sci. 1980 Apr 14;26(15): 1239-45.

Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins

This product is for Research Use Only· Not for Human or Veterinary or Therapeutic Use

Tel:781-999-4286 E_mail:info@targetmol.com Address:36 Washington Street,Wellesley Hills,MA 02481

Page 2 of 2 www.targetmol.com