# Data Sheet (Cat.No.T13127)



## Terbuthylazine

## **Chemical Properties**

CAS No.: 5915-41-3

Formula: C9H16ClN5

Molecular Weight: 229.71

Appearance: no data available

Storage: Powder: -20°C for 3 years | In solvent: -80°C for 1 year

## **Biological Description**

| Description   | Terbuthylazine is a triazine herbicide and a selective acetolactate synthase (ALS) inhibitor, toxic to aquatic organisms, and capable of disrupting gut microbiota. Terbuthylazine induces significant damage to the jejunal barrier and mitochondrial quality control imbalance.               |  |  |
|---------------|---|--|--|
| Targets(IC50) | Others  |  |  |
| In vitro      | Terbuthylazine induces innate immune response and inflammation by activating the cGAS-STING/NF-kB pathway in broiler heart muscle. [1] Terbuthylazine exposure activates the Nrf2/HO-1 pathway, leading to iron death. [2]  |  |  |
| In vivo       | Terbuthylazine (0.4 mg/kg) exposure caused cardiomyopathic rational changes in broilers. [1]  Terbuthylazine (0.4 or 4 mg/kg, 28 days) broilers showed pathological damage such as widening of interstitial space, breaking of muscle fibers, and deposition of myocardial collagen fibers. [2] |  |  |

## **Solubility Information**

| Solubility | DMSO: 40 mg/mL (174.13 mM), Sonication is recommended.          |  |
|------------|---|--|
|            | (< 1 mg/ml refers to the product slightly soluble or insoluble) |  |

### **Preparing Stock Solutions**

|       | 1mg       | 5mg        | 10mg       |
|-------|-----------|------------|------------|
| 1 mM  | 4.3533 mL | 21.7666 mL | 43.5332 mL |
| 5 mM  | 0.8707 mL | 4.3533 mL  | 8.7066 mL  |
| 10 mM | 0.4353 mL | 2.1767 mL  | 4.3533 mL  |
| 50 mM | 0.0871 mL | 0.4353 mL  | 0.8707 mL  |

Please select the appropriate solvent to prepare the stock solution, according to the solubility of the product in different solvents. Please use it as soon as possible.

Page 1 of 2 www.targetmol.com

#### Reference

Li H, et al. Terbuthylazine exposure induces innate immune response and inflammation through activating cGAS-STING/NF-κB pathway in myocardium of broiler chicken (Gallus gallus). Pestic Biochem Physiol. 2023 Dec;197: 105700.

Wu H, et al. Effects of terbuthylazine on myocardial oxidative stress and ferroptosis via Nrf2/HO-1 signaling pathway in broilers. Pestic Biochem Physiol. 2023 Dec;197:105698.

Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins

This product is for Research Use Only· Not for Human or Veterinary or Therapeutic Use

Tel:781-999-4286 E\_mail:info@targetmol.com Address:36 Washington Street,Wellesley Hills,MA 02481

Page 2 of 2 www.targetmol.com