


Hydrocortisone

Chemical Properties

CAS No. :	50-23-7
Formula:	C ₂₁ H ₃₀ O ₅
Molecular Weight:	362.46
Appearance:	no data available
Storage:	keep away from direct sunlight Powder: -20°C for 3 years In solvent: -80°C for 1 year



Biological Description

Description	Hydrocortisone (Cortisol) is a glucocorticoid hormone secreted by the adrenocortex. Hydrocortisone agonizes the glucocorticoid receptor, which promotes proteolytic metabolism, gluconeogenesis, capillary wall stabilization, and renal calcium excretion, as well as suppressing immune and inflammatory responses.
Targets(IC50)	Glucocorticoid Receptor, Endogenous Metabolite
In vitro	METHODS: Human articular chondrocytes were cultured in medium containing Hydrocortisone (0.05-1.0 µg/mL), and the relevant indexes were measured by Flow Cytometry. RESULTS: The catabolic signaling pathway of chondrocytes treated with 0.05 µg/mL Hydrocortisone was reduced, and the ability to synthesize ECM macromolecules was enhanced. [1] METHODS: Porcine brain capillary endothelial cells PBCEC were cultured with medium containing Hydrocortisone (550 nM) for seven days, and the cytoskeleton was examined using fluorescence microscopy. RESULTS: Hydrocortisone-induced cytoskeletal rearrangement. [2]
In vivo	In ex vivo guinea pig hearts, Hydrocortisone reduces post-ischemic oxidative stress, perfusion pressure, and effusion formation. Within the human cerebral microvascular endothelial cell line hCMEC/D3, Hydrocortisone prevents endothelial barrier disruption induced by pro-inflammatory stimulation (administration of TNFα), partly by maintaining occludin levels. Hydrocortisone induces class-switch recombination from Sµ to Sε in IL-4-treated B cells, supporting a model for sequential isotype transformation from IgM to IgE through IgG4. Following dendritic cell treatment, Hydrocortisone decreases the expression of MHC class II molecules, the dendritic-specific marker CD83, and costimulatory molecule CD86, while significantly reducing IL-12 secretion. Additionally, it inhibits the increase of IL-4 (induced by IFN-γ) but does not affect IL-5. Hydrocortisone lowers T cell proliferation in dendritic cells. It prevents TNF-α-induced severe degradation of glycocalyx, increases in coronary resistance, vascular leakage and permeability, hydroxyethyl starch, and causes degranulation of mast cells in ex vivo guinea pig hearts. Hydrocortisone inhibits the shedding of multiligand glycosaminoglycans-1, heparan sulfate, and hyaluronic acid post-ischemia and suppresses the release of histamine from resident mast cells.

A DRUG SCREENING EXPERT

Cell Research	Cells are plated on top of collagen IV-coated transwell chambers for six-well plates (24 mm diameter, membrane material: polyethylene terephthalate (PET), 0.4 μ m pores, pore density 1.6 $\times$ 10 ⁶ ?cm ²) at densities of 2.5 $\times$ 10 ⁴ ?cells cm ² ?per well. When they have reached confluence at day 5, the different experimental sets of cells are transferred to differentiation medium containing reduced amounts of FCS and treated with TNF α or hydrocortisone as indicated.
---------------	--

Solubility Information

Solubility	DMSO: 50 mg/mL (137.95 mM),Sonication is recommended. Ethanol: 22 mg/mL (60.7 mM),Sonication is recommended. 10% DMSO+40% PEG300+5% Tween 80+45% Saline: 6.8 mg/mL (18.76 mM),Solution. (< 1 mg/ml refers to the product slightly soluble or insoluble)
------------	--

Preparing Stock Solutions

	1mg	5mg	10mg
1 mM	2.7589 mL	13.7946 mL	27.5893 mL
5 mM	0.5518 mL	2.7589 mL	5.5179 mL
10 mM	0.2759 mL	1.3795 mL	2.7589 mL
50 mM	0.0552 mL	0.2759 mL	0.5518 mL

Please select the appropriate solvent to prepare the stock solution, according to the solubility of the product in different solvents. Please use it as soon as possible.

Reference

- Wang J, et al. Physiological levels of hydrocortisone maintain an optimal chondrocyte extracellular matrix metabolism. *Ann Rheum Dis*. 2004 Jan;63(1):61-6.
- Lei H, Wang Z, Jiang D, et al. CRISPR screening identifies CDK12 as a conservative vulnerability of prostate cancer. *Cell Death & Disease*. 2021, 12(8): 1-11.
- Schrot S, et al. Influence of hydrocortisone on the mechanical properties of the cerebral endothelium in vitro. *Biophys J*. 2005 Dec;89(6):3904-10.
- Li J, Bai Y, Liu Y, et al. Transcriptome-based chemical screens identify CDK8 as a common barrier in multiple cell reprogramming systems. *Cell Reports*. 2023, 42(6).
- Ji X, Jiang P, Li Y, et al. Maternal Bisphenol B Exposure and Mammary Gland Development of Offspring: A Time-Series Analysis. *Environment & Health*. 2023
- Bergt S, et al. Hydrocortisone reduces the beneficial effects of toll-like receptor 2 deficiency on survival in a mouse model of polymicrobial sepsis. *Shock*. 2013 Nov;40(5):414-9.
- Li Y, et al. Risk of death does not alter the efficacy of hydrocortisone therapy in a mouse E. coli pneumonia model: risk and corticosteroids in sepsis. *Intensive Care Med*. 2008 Mar;34(3):568-77.
- Zhang D, Ma B, Liu D, et al. Discovery of a peptide proteolysis-targeting chimera (PROTAC) drug of p300 for prostate cancer therapy. *EBioMedicine*. 2024, 105.
- Liang L, Shen Z, Zhang K, et al. DLGAP5 upregulates E2F1 to promote prostate adenocarcinoma neuroendocrine differentiation. *Life Sciences*. 2025: 123442.
- Jabara HH, et al. *J Immunol*, 1993, 151(9), 4528-4533.

Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins

This product is for Research Use Only · Not for Human or Veterinary or Therapeutic Use

Tel: 781-999-4286 E_mail: info@targetmol.com Address: 36 Washington Street, Wellesley Hills, MA 02481