


Afatinib Dimaleate

Chemical Properties

CAS No. :	850140-73-7
Formula:	C32H33ClFN5O11
Molecular Weight:	718.08
Appearance:	no data available
Storage:	Powder: -20°C for 3 years In solvent: -80°C for 1 year



Biological Description

Description	Afatinib Dimaleate (BIBW 2992MA2) is an orally bioavailable anilino-quinazoline derivative and inhibitor of the receptor tyrosine kinase (RTK) epidermal growth factor receptor (ErbB; EGFR) family, with antineoplastic activity.
Targets(IC50)	EGFR,HER,Autophagy
In vitro	In the MDA-MB-453 transplantation tumor model, oral administration of Afatinib (20 mg/kg), which down-regulated the phosphorylation levels of EGFR and AKT, induced tumor regression. In the NCI-N87 transplantation model of HER2-positive gastric cancer, oral administration of Afatinib (25 mg/kg) was able to eliminate tumors. In A7, A431, FaDu, UT-SCC-14 and UT-SCC-15 transplantation tumor models, oral administration of Afatinib (30 mg/kg) inhibited tumor growth.
In vivo	In lung cancer cell lines expressing wild-type (H1666) or L858R/T790M (NCI-H1975) EGFR, Afatinib inhibited cell growth more effectively. In NSCLC cell lines expressing HER2 776insV (NCI-H1781) or EGFR E746_A750del (HCC827), Afatinib was more effective in inhibiting cell growth.
Kinase Assay	In vitro kinase activity assay: EGFR kinase: Each 100 μ L enzyme reaction contained 10 μ L of inhibitor in 50% Me2SO, 20 μ L of substrate solution (200 mM HEPES pH 7.4, 50 mM Mg-acetate, 2.5 mg/mL poly (EY), 5 μ g/mL bio-pEY) and 20 μ L enzyme preparation. The enzymatic reaction is started by addition of 50 μ L of a 100 μ M ATP solution made in 10 mM MgCl2. Assays are carried out at room temperature for 30 min and terminated by the addition of 50 μ L of stop solution (250 mM EDTA in 20 mM HEPES pH 7.4). 100 μ L are transferred to a streptavidin coated microtiterplate, after an incubation time of 60 min at room temperature the plate is washed with 200 μ L of wash solution (50 mM Tris, 0.05% Tween20). A 100 μ L aliquot of a HRPO- labeled anti-PY antibody (PY20H Anti-Ptyr:HRP) 250 ng/mL are added to the wells. After 60 min of incubation, the plate is washed three times with a 200 μ L wash solution. The samples are then developed with a 100 μ L TMB Peroxidase Solution (A:B= 1:1). The reaction is stopped after 10 min. The plate is transferred to an ELISA reader and extinction is measured at OD450 nM. HER2-IC enzyme: Enzyme activity is assayed in the presence or absence of serial inhibitor dilutions performed in 50 % Me2SO. Each 100 μ L reaction contains similar components as described for EGFR kinase assay with addition of 1000 μ M Na3VO4. The enzymatic reaction is started by addition of 50 μ L of 500 μ M ATP solution made in 10 mM Mg-acetate. The dilution of the enzyme is set so that incorporation of phosphate into bio-pEY is linear with respect to time and amount of enzyme. The enzyme preparation is

diluted in 20 mM HEPES pH 7.4, 130 mM NaCl, 0.05% Triton X-100, 1 mM DTT and 10% glycerol. Assays are carried out at room temperature for 30 min and terminated by the addition of 50 μ L of stop solution. Src kinase assays: Each 100 μ L reaction contained 10 μ L of inhibitor in 50 % Me₂SO, 20 μ L of enzyme preparation, 20 μ L of substrate solution supplemented with 1000 μ M Na₃VO₄. The enzymatic reaction is started by addition of 50 μ L of a 1000 μ M ATP solution made in 10 mM Mg-acetate. BIRK kinase assay: 250 mM Tris pH 7.4, 10 mM DTT, 2.5 mg/mL poly(EY), 5 mg/mL bio-pEY is used as substrate solution and enzymatic reaction is started by addition of 50 μ L of a 2 mM ATP solution made in 8 mM MnCl₂, 20 mM Mg-acetate. VEGF2 and HGFR kinase assays: Assays are carried out at room temperature for 20 minutes and terminated by the addition of 10 μ L of 5 % H₃PO₄. The precipitate is then trapped onto GF/B filters using a 96 well filter mate universal harvester. After extensive washing the filter plate is dried for 1 h at 50°C, sealed and incorporated radioactivity is determined by scintillation counting using a TopCount[®] or a Microbeta b counter[®].

Cell Research

Cytotoxicity is determined using MTT assay. The IC₅₀ value is defined as the drug concentration resulting in 50% cell death. Both the fitted sigmoidal dose response curve and IC₅₀ are calculated by Bliss method.(Only for Reference)

Solubility Information

Solubility

H₂O: < 1 mg/mL (insoluble or slightly soluble),
DMSO: 50 mg/mL (69.63 mM), Sonication is recommended.
Ethanol: < 1 mg/mL (insoluble or slightly soluble),
(< 1 mg/mL refers to the product slightly soluble or insoluble)

Preparing Stock Solutions

	1mg	5mg	10mg
1 mM	1.3926 mL	6.963 mL	13.926 mL
5 mM	0.2785 mL	1.3926 mL	2.7852 mL
10 mM	0.1393 mL	0.6963 mL	1.3926 mL
50 mM	0.0279 mL	0.1393 mL	0.2785 mL

Please select the appropriate solvent to prepare the stock solution, according to the solubility of the product in different solvents. Please use it as soon as possible.

Reference

D Li, et al. Oncogene. 2008, 27(34):4702-4711.

Li C, Sun Y D, Yu G Y, et al. Integrated Omics of Metastatic Colorectal Cancer. Cancer Cell. 2020, 38(5): 734-747. e9

Mizuma K, Takashima A, Cubitt B, et al. The Pan-ErbB tyrosine kinase inhibitor afatinib inhibits multiple steps of the mammarenavirus life cycle. Virology. 2022

Wang XK, et al. Cancer Res. 2014, 74(16):4431-4445.

Gurtner K, et al. Radiat Oncol. 2014, 9:261.

Schwab CL, et al. Br J Cancer. 2014, 111(9):1750-1756.

Janjiqian YY, et al. J Nucl Med. 2013, 54(6):936-943.

Li C, Sun Y D, Yu G Y, et al. Integrated Omics of Metastatic Colorectal Cancer[J]. Cancer Cell. 2020, 38(5): 734-747. e9

Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins

This product is for Research Use Only· Not for Human or Veterinary or Therapeutic Use

Tel:781-999-4286 E_mail:info@targetmol.com Address:36 Washington Street,Wellesley Hills,MA 02481