


Capivasertib

Chemical Properties

CAS No. :	1143532-39-1
Formula:	C ₂₁ H ₂₅ ClN ₆ O ₂
Molecular Weight:	428.92
Appearance:	no data available
Storage:	Powder: -20°C for 3 years In solvent: -80°C for 1 year



Biological Description

Description	Capivasertib (AZD5363) is a broad-spectrum AKT inhibitor with inhibitory activity against Akt1, Akt2, and Akt3 (IC ₅₀ =3/7/7 nM) with oral activity. Capivasertib has antitumor activity for the treatment of breast cancer.
Targets(IC ₅₀)	Akt, Autophagy, mTOR, PKA
In vitro	METHODS: Six strains of human gastric cancer cells were treated with Capivasertib (40 nM-50 μM) for 72 h. Cell viability was measured by SRB Assay. RESULTS: The IC₅₀ of Capivasertib on HGS27, AGS, N87, SNU-1, MKN45, and MGC803 cells were 4.6/0.1/14.18/24.04/30.0/44.4 μM, respectively. [1] METHODS: Breast cancer cells BT474c and prostate cancer cells LNCaP were treated with Capivasertib (0.03-10 μmol/L) for 2 h, and the expression levels of target proteins were detected by Western Blot. RESULTS: Capivasertib effectively inhibited the phosphorylation of S6 and 4E-BP1 in the cell lines, while it increased the phosphorylation of AKT at ser473 and thr308. [2]
In vivo	METHODS: To assay antitumor activity in vivo, Capivasertib (100-300 mg/kg, 10% DMSO 25% w/v Kleptose HPB buffer) was administered by gavage twice daily for two weeks to nude mice harboring mammary cancer tumor BT474c. RESULTS: Capivasertib dose-dependently inhibited the growth of human tumor xenografts in vivo. [2] METHODS: To assay antitumor activity in vivo, Capivasertib (100-300 mg/kg, 10% DMSO 25% w/v Kleptose HPB buffer) was administered by gavage to a PDGfCX mouse model twice daily for twenty days. RESULTS: Capivasertib monotherapy resulted in 60% tumor growth inhibition. [3]
Kinase Assay	The ability of AZD5363 to inhibit the activity of AKT1, AKT2 and AKT3 was evaluated by the Caliper Off-Chip Incubation Mobility Shift Assay. Active recombinant AKT1, AKT2, or AKT3 were incubated with a 5-FAM-labeled custom-synthesized peptide substrate together with increasing concentrations of inhibitor. Final reactions contained 1 to 3 nmol/L AKT1, AKT2, or AKT3 enzymes; 1.5 μmol/L peptide substrate; ATP at Km for each AKT isoform; 10 mmol/L MgCl ₂ , 4 mmol/L dithiothreitol (DTT), 100 mmol/L HEPES, and 0.015% Brij-35. The reactions were incubated at room temperature for 1 hour and stopped by the addition of a buffer containing 100 mmol/L HEPES, 0.015% Brij-35 solution, 0.1% coating reagent, 40 mmol/L EDTA, and 5% DMSO. Plates were then analyzed using a Caliper LC3000, allowing for separation of peptide substrate and phosphorylated product by electrophoresis with subsequent detection and

quantification of laser-induced fluorescence. To determine the kinase selectivity profile, AZD5363 was also tested against PKA, ROCK1, ROCK2, and P70S6K. PKA, ROCK1, and ROCK2 activity were determined by Caliper Off-Chip Incubation Mobility Shift Assay, as described earlier. Final reaction conditions for measuring ROCK1 activity were 5 nmol/L active recombinant ROCK1, 1.5 μ mol/L fluorescein isothiocyanate (FITC)-labeled custom peptide substrate, 7 μ mol/L ATP, 1 mmol/L DTT, 5 mmol/L MgCl₂, 100 mmol/L HEPES, 0.015% Brij-35, and 5 mmol/L β -glycerophosphate; final reaction for measuring ROCK2 activity contained 7.5 nmol/L active recombinant ROCK2, 1.5 μ mol/L FAM-labeled custom peptide substrate, 7.5 μ mol/L ATP, 1 mmol/L DTT, 10 mmol/L MgCl₂, 100 mmol/L HEPES, 0.015% Brij-35, and 5 mmol/L β -glycerophosphate; and protein kinase A (PKA) activity was measured in a final reaction containing 0.0625 nmol/L PKA, 3 μ mol/L FITC-labeled custom peptide substrate, 4.6 μ mol/L ATP, 1 mmol/L DTT, 10 mmol/L MgCl₂, 110 mmol/L HEPES, and 0.015% Brij-35. P70S6K activity was measured by a radioactive (³³P-ATP) filter-binding assay. Recombinant S6K1 (T412E) was assayed against a substrate peptide (KKRNRTLTV) in a final volume of 25.5 μ L containing 8 mmol/L MOPS, 200 μ mol/L EDTA, 100 μ mol/L substrate peptide, 10 mmol/L magnesium acetate, 20 μ mol/L γ -³³P-ATP (50–1,000 cpm/pmol), and increasing concentrations of AZD5363. The reactions were incubated for 30 minutes at room temperature and terminated by the addition of 0.5 mol (3%) orthophosphoric acid. Reactions were then harvested onto a P81 UniFilter and product formation quantified. IC₅₀ values for all enzyme assays were obtained by fitting data in Origin 7.0.

Cell Research	A high-throughput screening cell-based assay was developed to measure cellular AKT activity using the MDA-MB-468 breast cancer cell line. Cells were exposed to AZD5363 at concentrations ranging from 3 to 0.003 μ mol/L. After a 2-hour treatment, cells were fixed with formaldehyde, washed, permeabilized with 0.5% polysorbate 20 and then probed with a phospho-specific antibody against GSK3 β ser9. Levels of phosphorylated GSK3 β ser9 were measured with an Acumen Explorer laser scanning cytometer and IC ₅₀ values estimates by fitting data in Origin 7.0.
Animal Research	When mean tumor sizes reached approximately 0.2 cm ³ , the mice were randomized into control and treatment groups. The treatment groups received varying dose schedules of AZD5363 solubilized in a 10% DMSO 25% w/v Kleptose HPB buffer by oral gavage, docetaxel solubilized in 2.6% ethanol in injectable water by intravenous injection once on day 1 at 15 or 5 mg/kg once weekly. When administered in combination, docetaxel was administered 1 hour before the oral dose of AZD5363. The control group received the DMSO/Kleptose buffer alone, twice daily by oral gavage. Tumor volumes (measured by caliper), animal body weight, and tumor condition were recorded twice weekly for the duration of the study. Mice were sacrificed by CO ₂ euthanasia. The tumor volume was calculated (taking length to be the longest diameter across the tumor and width to be the corresponding perpendicular diameter) using the formula: (length $\times$ width) $\times \sqrt{(\text{length} \times \text{width}) \times (\pi/6)}$. Growth inhibition from the start of treatment was assessed by comparison of the differences in tumor volume between control and treated groups. Because the variance in mean tumor volume data increases proportionally with volume (and is therefore disproportionate between groups), data were log transformed to remove any size dependency before statistical evaluation. Statistical significance was evaluated using a one-tailed, 2-sample t-test.

Solubility Information

Solubility	Ethanol: 1 mg/mL (2.33 mM), Sonication is recommended. DMSO: 50 mg/mL (116.57 mM), Sonication is recommended. 10% DMSO+40% PEG300+5% Tween 80+45% Saline: 8 mg/mL (18.65 mM), Solution.
------------	---

H₂O: Insoluble,
(< 1 mg/ml refers to the product slightly soluble or insoluble)

Preparing Stock Solutions

	1mg	5mg	10mg
1 mM	2.3314 mL	11.6572 mL	23.3144 mL
5 mM	0.4663 mL	2.3314 mL	4.6629 mL
10 mM	0.2331 mL	1.1657 mL	2.3314 mL
50 mM	0.0466 mL	0.2331 mL	0.4663 mL

Please select the appropriate solvent to prepare the stock solution, according to the solubility of the product in different solvents. Please use it as soon as possible.

Reference

- Wang J, et al. Ceritinib increases sensitivity of AKT inhibitors to gastric cancer. *Eur J Pharmacol.* 2021 Apr 5;896: 173879.
- Weng Q, Zhao M, Zheng J, et al. STAT3 dictates β -cell apoptosis by modulating PTEN in streptozocin-induced hyperglycemia. *Cell Death & Differentiation.* 2020, 27(1): 130-145
- Davies BR, et al. Preclinical pharmacology of AZD5363, an inhibitor of AKT: pharmacodynamics, antitumor activity, and correlation of monotherapy activity with genetic background. *Mol Cancer Ther.* 2012 Apr;11(4):873-87.
- Liu Q, Miao Y, Wang X, et al. Structure-based virtual screening and biological evaluation of novel non-bisphosphonate farnesyl pyrophosphate synthase inhibitors. *European Journal of Medicinal Chemistry.* 2019: 111905
- Wang J, Xu X, Wang T, et al. Ceritinib increases sensitivity of AKT inhibitors to gastric cancer. *European Journal of Pharmacology.* 2021: 173879.
- Li J, et al. The AKT inhibitor AZD5363 is selectively active in PI3KCA mutant gastric cancer, and sensitizes a patient-derived gastric cancer xenograft model with PTEN loss to Taxotere. *J Transl Med.* 2013 Oct 2;11:241.
- Weng Q, Zhao M, Zheng J, et al. STAT3 dictates β -cell apoptosis by modulating PTEN in streptozocin-induced hyperglycemia[J]. *Cell Death & Differentiation.* 2020, 27(1): 130-145.
- Wang J, Xu X, Wang T, et al. Ceritinib increases sensitivity of AKT inhibitors to gastric cancer. *European Journal of Pharmacology.* 2021: 173879.
- Bing S, Xiang S, Xia Z, et al. AKT inhibitor Hu7691 induces differentiation of neuroblastoma cells. *Acta Pharmaceutica Sinica B.* 2023
- Zhang W, Li X, Jiang M, et al. SOCS3 deficiency-dependent autophagy repression promote the survival of early-stage myeloid-derived suppressor cells in breast cancer by activating the Wnt/mTOR pathway. *Journal of Leukocyte Biology.* 2023: qiad020.
- Zhang K M, Zhao D C, Li Z Y, et al. Inactivated cGAS-STING Signaling Facilitates Endocrine Resistance by Forming a Positive Feedback Loop with AKT Kinase in ER+ HER2-Breast Cancer. *Advanced Science.* 2024: 2403592.

Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins

This product is for Research Use Only · Not for Human or Veterinary or Therapeutic Use

Tel: 781-999-4286 E_mail: info@targetmol.com Address: 36 Washington Street, Wellesley Hills, MA 02481