

c-Kit-IN-3 D-tartrate

Chemical Properties


CAS No. :

Formula: C30H26ClF3N2O10

Molecular Weight: 666.98

Appearance: no data available

Storage: Powder: -20°C for 3 years | In solvent: -80°C for 1 year



Biological Description

Description	c-Kit-IN-3 D-tartrate is a potent and selective c-KIT kinase inhibitor (IC50s: 4 nM, 8 nM for c-KIT wt, and c-KIT T670I).
Targets(IC50)	Others
In vitro	c-Kit-IN-3 D-tartrate (Compound 18; 0.1-10 μ M; 6 days; primary GIST patient cells) exhibits dose-dependent antiproliferative effects. c-Kit-IN-3 (0.01-1 μ M; 24 hours; GIST-T1, GIST-T1-T670I, and GIST-5R cells) treatment induces dose-dependent cell apoptotic death. c-Kit-IN-3 (0.01-1 μ M; 24 hours; GIST-T1, GIST-T1-T670I, and GIST-5R cells) treatment arrests the cell cycle into the G0/G1 phase in all of these three cell lines. c-Kit-IN-3 (0-1 μ M; 24 hours) blocks the autophosphorylation of c-KIT pY703, pY719, and pY823 in GIST-T1, GIST-T1-T670I, and GIST-5R, respectively, cells at a concentration of 30 nM and inhibits the downstream signaling mediators pAKT (T308/S473), pS6 (S235/236), and pERK (T202/204). c-Kit-IN-3 potently inhibits the activity of CSF1R (IC50: 18 nM), PDGFR α (IC50: 25 nM), RET (IC50: 34 nM), and it relatively less potently inhibits DDR1 (IC50: 135 nM), FLT4 (IC50: 121 nM), and PDGFR β (IC50: 97 nM). c-Kit-IN-3 (0.006 μ M-1.37 μ M) potently inhibits the growth of c-KIT-dependent GIST cancer cells, such as GIST-T1 (IC50: 0.006 μ M); GIST-882 (IC50: 0.013 μ M); GIST-T1-T670I (IC50 : 0.011 μ M); GIST-5R (IC50: 0.073 μ M); GIST-48B (IC50: 1.37 μ M), respectively.
In vivo	c-Kit-IN-3 D-tartrate (oral gavage; 20-100 mg/kg/day; for 11 days; female BALB/C-nu mice) treatment dose-dependently inhibits the tumor progression. c-Kit-IN-3 (1 mg/kg iv for mice, rats, dog; 10 mg/kg p.o. for mice, rats; and 5 mg/kg p.o. for dog) has T1/2 of 4.5 h, 6.4 h, 19.4 h for mice, rats and dogs, respectively. And it possesses acceptable bioavailability in mice (F = 43%), rats (F = 50%), and dogs (F = 81%).

Preparing Stock Solutions

	1mg	5mg	10mg
1 mM	1.4993 mL	7.4965 mL	14.993 mL
5 mM	0.2999 mL	1.4993 mL	2.9986 mL
10 mM	0.1499 mL	0.7496 mL	1.4993 mL
50 mM	0.030 mL	0.1499 mL	0.2999 mL

Please select the appropriate solvent to prepare the stock solution, according to the solubility of the product in different solvents. Please use it as soon as possible.

Reference

Wu Y, et al. Discovery of 2-(4-Chloro-3-(trifluoromethyl)phenyl)-N-(4-((6,7-dimethoxyquinolin-4-yl)oxy)phenyl) acetamide (CHMFL-KIT-64) as a Novel Orally Available Potent Inhibitor against Broad-Spectrum Mutants of c-KIT Kinase for Gastrointestinal Stromal Tumors. J Med Chem. 2019 Jul 11;62(13):6083-6101.

Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins

This product is for Research Use Only. Not for Human or Veterinary or Therapeutic Use

Tel: 781-999-4286 E_mail: info@targetmol.com Address: 36 Washington Street, Wellesley Hills, MA 02481