

Spautin-1

Chemical Properties


CAS No. : 1262888-28-7

Formula: C₁₅H₁₁F₂N₃

Molecular Weight: 271.26

Appearance: no data available

Storage: Powder: -20°C for 3 years | In solvent: -80°C for 1 year



Biological Description

Description	Spautin-1 is an autophagy inhibitor that suppresses the deubiquitination activity of ubiquitin-specific peptidase USP10 and USP13. Spautin-1 inhibits UPR and exhibits preferential cytotoxicity in glucose-starved cancer cells.
Targets(IC50)	Apoptosis, Autophagy, DUB
In vitro	In Bcap-37 cells, Spautin-1 dramatically enhanced cell death in glucose-free media and induces apoptotic morphology. In Bax-Bak DKO cells, spautin-1 inhibits etoposide induced autophagic cell death. Spautin-1 promotes the degradation of Vps34 complexes by regulating the deubiquitination activity of USP10 and USP13, and reduces the levels of PtdIns3P. [1] In PDGF-treated cells, spautin-1 stabilizes α -smooth muscle cell actin and calponin, prevents actin filament disorganization, diminishes production of extracellular matrix, and abrogates VSMC hyperproliferation and migration. [2] In CML cells, spautin-1 markedly inhibits IM-induced autophagy by downregulating Beclin-1, and enhances IM-induced apoptosis by inactivating PI3K/AKT and activating downstream GSK3 β . [3] Spautin-1 also specifically reduces infectious dengue virus titers in BHK-21 cells. [4]
In vivo	Spautin-1 effectively mitigates the pathogenesis of acute pancreatitis induced by cerulein or L-arginine. Pretreatment with Spautin-1 significantly reduces elevated serum amylase and lipase levels, markers of trypsin activity, and inhibits the increase of serum TNF α caused by cerulein. Additionally, Spautin-1 treatment alleviates inflammation-induced damage such as edema, degeneration, coagulative necrosis, and the infiltration of inflammatory cells[5].
Kinase Assay	Kinase inhibitory assays in Vitro: The inhibitory ability against each kinase except for MEK1 and Raf-1 is evaluated by examining their ability to phosphorylate various substrate peptides in the presence of CH5424802 using time-resolved fluorescence resonance energy transfer (TR-FRET) assay or fluorescence polarization (FP) assay. The inhibitory activity against MEK1 is evaluated by quantitative analysis of the phosphorylation of a substrate peptide by a recombinant ERK2 protein in the presence of CH5424802. The inhibitory activity against Raf-1 is evaluated by examining the ability of the kinases to phosphorylate MEK1 in the presence of CH5424802.
Cell Research	Spautin-1 is dissolved in DMSO. Cell proliferation is evaluated using CCK-8 kit. K562 cells (1x10 ⁵ /mL) are seeded into 96-well plates in triplicate and then treated with 125 to 4,000 nM IM alone or in combination with spautin-1 (10 μ M). After 48 h of incubation, 10 μ L of CCK-8 reagent is added to each well. Four hours later, the absorbance is read at

450 nm using a microplate reader[1].

Solubility Information

Solubility	DMSO: 45 mg/mL (165.89 mM), Sonication is recommended. (< 1 mg/ml refers to the product slightly soluble or insoluble)
------------	---

Preparing Stock Solutions

	1mg	5mg	10mg
1 mM	3.6865 mL	18.4325 mL	36.865 mL
5 mM	0.7373 mL	3.6865 mL	7.373 mL
10 mM	0.3687 mL	1.8433 mL	3.6865 mL
50 mM	0.0737 mL	0.3687 mL	0.7373 mL

Please select the appropriate solvent to prepare the stock solution, according to the solubility of the product in different solvents. Please use it as soon as possible.

Reference

- Guo J, et al. Potent USP10/13 antagonist spautin-1 suppresses melanoma growth via ROS-mediated DNA damage and exhibits synergy with cisplatin. *J Cell Mol Med*. 2020 Apr; 24(7):4324-4340. doi: 10.1111/jcmm.15093. Epub 2020 Mar 4.
- Wu W, Xu H, Liao C, et al. Blockade of USP14 potentiates type I interferon signaling and radiation-induced antitumor immunity via preventing IRF3 deubiquitination. *Cellular Oncology*. 2022: 1-15
- Shao S, et al. Spautin-1, a novel autophagy inhibitor, enhances imatinib-induced apoptosis in chronic myeloid leukemia. *Int J Oncol*. 2014 May; 44(5):1661-8.
- Yuan T, Chen Z, Yan F, et al. Deubiquitinating Enzyme USP10 Promotes Hepatocellular Carcinoma Metastasis through Deubiquitinating and Stabilizing Smad4 Protein. *Molecular Oncology*. 2020, 14(1): 197-210
- Yue X, Liu T, Wang X, et al. Pharmacological inhibition of BAP1 recruits HERC2 to competitively dissociate BRCA1-BARD1, suppresses DNA repair and sensitizes CRC to radiotherapy. *Acta Pharmaceutica Sinica B*. 2023
- Xiao J, et al. Spautin-1 Ameliorates Acute Pancreatitis via Inhibiting Impaired Autophagy and Alleviating Calcium Overload. *Mol Med*. 2016 Oct; 22:643-652.
- Xu X, Xie T, Zhou M, et al. Hsc70 promotes anti-tumor immunity by targeting PD-L1 for lysosomal degradation. *Nature Communications*. 2024, 15(1): 4237.
- Zhang LM, et al. Spautin-1 administration mitigates mild TBI-induced cognitive and memory dysfunction in mice via activation of caspase-3. *Int Immunopharmacol*. 2023 Apr; 117:109906.
- Xiao J, et al. Spautin-1 Ameliorates Acute Pancreatitis via Inhibiting Impaired Autophagy and Alleviating Calcium Overload. *Mol Med*. 2016 Aug 18;22.
- Yuan T, Chen Z, Yan F, et al. Deubiquitinating Enzyme USP10 Promotes Hepatocellular Carcinoma Metastasis through Deubiquitinating and Stabilizing Smad4 Protein[J]. *Molecular oncology*. 2020, 14(1): 197-210

Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins

This product is for Research Use Only. Not for Human or Veterinary or Therapeutic Use

Tel: 781-999-4286 E_mail: info@targetmol.com Address: 36 Washington Street, Wellesley Hills, MA 02481