# Data Sheet (Cat.No.T1968)



#### UNC2250

Storage:

## **Chemical Properties**

CAS No.: 1493694-70-4


Formula: C24H36N6O2

Molecular Weight: 440.58

Appearance: no data available

store under nitrogen

Powder: -20°C for 3 years | In solvent: -80°C for 1 year



# **Biological Description**

Description	UNC2250 is an effective and specific Mer inhibitor (IC50=1.7 nM).			
Targets(IC50)	TAM Receptor			
In vivo	According to in vivo PK experiment, UNC2250 has a moderate half-life, clearance, and volume of distribution as well as reasonable oral bioavailability and good solubility. [1]			
Kinase Assay	Microfluidic Capillary Electrophoresis (MCE) Assay: Activity assays are performed in a 384-well, polypropylene microplate in a final volume of 50 $\mu$ L of 50 mM Hepes, pH 7.4, containing 10 mM MgCl2, 1.0 mM DTT, 0.01% Triton X-100, 0.1% bovine serum albumin (BSA), containing 1.0 $\mu$ M fluorescent substrate and ATP at the Km for each enzyme. All reactions are terminated by addition of 20 $\mu$ L of 70 mM EDTA. After an 180 min incubation, phosphorylated and unphosphorylated substrate peptides are separated in buffer supplemented with 1× CR-8 on a LabChip EZ Reader equipped with a 12-sipper chip. Data are analyzed using EZ Reader software.			
Cell Research	BT-12 rhabdoid tumor cells (10?000 cells) are cultured in 2.0 mL of 0.35% soft agar containing 0.5× RPMI medium, 7.5% FBS, and the indicated concentrations of 10 or DMSO vehicle only and overlaid with 0.5 mL of 1× RPMI medium containing 15% FBS and 10 or DMSO vehicle only. Medium and 10 or vehicle are refreshed 2 times per week. Colonies were stained with thiazolyl blue tetrazolium bromide and counted after 3 weeks. Colo699 NSCLC cells (15?000 cells) are cultured in 1.5 mL of 0.35% soft agar containing 1× RPMI medium and 10% FBS and overlaid with 2.0 mL of 1× RPMI medium containing 10% FBS and the indicated concentrations of 10 or DMSO vehicle only. Medium and 10 or vehicle are refreshed 3 times per week. Colonies are stained with nitrotetrazolium blue chloride and counted after 2 weeks.(Only for Reference)			

# **Solubility Information**

Solubility	DMSO: 25 mg/mL (56.74 mM),Sonication is recommended.
	(< 1 mg/ml refers to the product slightly soluble or insoluble)

Page 1 of 2 www.targetmol.com

## **Preparing Stock Solutions**

	1mg	5mg	10mg
1 mM	2.2697 mL	11.3487 mL	22.6974 mL
5 mM	0.4539 mL	2.2697 mL	4.5395 mL
10 mM	0.227 mL	1.1349 mL	2.2697 mL
50 mM	0.0454 mL	0.227 mL	0.4539 mL

Please select the appropriate solvent to prepare the stock solution, according to the solubility of the product in different solvents. Please use it as soon as possible.

#### Reference

Zhang W, et al. J Med Chem. 2013, 56(23), 9683-9692.

Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins

This product is for Research Use Only· Not for Human or Veterinary or Therapeutic Use

Tel:781-999-4286 E\_mail:info@targetmol.com Address:36 Washington Street,Wellesley Hills,MA 02481

Page 2 of 2 www.targetmol.com