


Articaine hydrochloride

Chemical Properties

CAS No. :	23964-57-0
Formula:	C ₁₃ H ₂₁ ClN ₂ O ₃ S
Molecular Weight:	320.84
Appearance:	no data available
Storage:	Powder: -20°C for 3 years In solvent: -80°C for 1 year



Biological Description

Description	Articaine hydrochloride (Hoe-045), a thiophene-containing local anesthetic, is pharmacologically similar to mepivacaine.
Targets(IC50)	Sodium Channel
In vivo	The time to maximum drug concentrations of articaine occurs about 10 to 15 minutes after submucosal injection of articaine 4% 80 mg. The elimination half-time of articaine is about 20 minutes. Articaine is better able to diffuse through soft tissue and bone than other local anaesthetics, the concentration of articaine in the alveolus of a tooth in the upper jaw after extraction is about 100 times higher than that in systemic circulation. [1] Articaine: VAS (Visual Analogue Scale) scores (from 0 to 10 cm) by patients 4 to < 13 years of age are 0.5 for simple procedures and 1.1 for complex procedures, and average investigator scores are 0.4 and 0.6 for simple and complex procedures, respectively. No serious adverse events related to the articaine occurs, the only adverse event considered related to articaine is accidental lip injury in one patient. [2] Articaine results in success rate of 64.5% in electronic pulp testing in healthy adult volunteers injected with 4% articaine. Articaine infiltration produces significantly more episodes of no response to maximum stimulation in first molars than lidocaine. Mandibular buccal infiltration is more effective with 4% articaine with epinephrine compared to 2% lidocaine with epinephrine. [3] Articaine (4%) results in the success rate of 24% for the inferior alveolar nerve block in randomized, double-blind study. [4] Articaine formulation results in successful pulpal anesthesia ranged from 75 to 92 percent and onset of pulpal anesthesia ranged from 4.2 to 4.7 minutes in healthy volunteer. For articaine, 4 percent (two of 56) of the subjects reported swelling and no subjects reported bruising. Ninety-eight percent (59 of 60) of the subjects had lip numbness with the articaine solution. [5]

Solubility Information

Solubility	DMSO: 60 mg/mL (187.01 mM),Sonication is recommended. H2O: 59 mg/mL (183.89 mM),Sonication is recommended. Ethanol: 60 mg/mL (187.01 mM),Sonication is recommended. (< 1 mg/ml refers to the product slightly soluble or insoluble)
------------	--

A DRUG SCREENING EXPERT

Preparing Stock Solutions

	1mg	5mg	10mg
1 mM	3.1168 mL	15.5841 mL	31.1682 mL
5 mM	0.6234 mL	3.1168 mL	6.2336 mL
10 mM	0.3117 mL	1.5584 mL	3.1168 mL
50 mM	0.0623 mL	0.3117 mL	0.6234 mL

Please select the appropriate solvent to prepare the stock solution, according to the solubility of the product in different solvents. Please use it as soon as possible.

Reference

Oertel R, et al. Clin Pharmacokinet, 1997, 33(6), 417-425.

Malamed SF, et al. Pediatr Dent, 2000, 22(4), 307-311.

Kanaa MD, et al. J Endod, 2006, 32(4), 296-298.

Claffey E, et al. J Endod, 2004, 30(8), 568-571.

Robertson D, et al. J Am Dent Assoc, 2007, 138(8), 1104-1112.

Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins

This product is for Research Use Only · Not for Human or Veterinary or Therapeutic Use

Tel:781-999-4286 E_mail:info@targetmol.com Address:36 Washington Street,Wellesley Hills,MA 02481