

## Bergapten

## Chemical Properties

CAS No. : 484-20-8


Formula: C<sub>12</sub>H<sub>8</sub>O<sub>4</sub>

Molecular Weight: 216.19

Appearance: no data available

Storage: keep away from direct sunlight

Powder: -20°C for 3 years | In solvent: -80°C for 1 year



## Biological Description

Description	Bergapten (5-Methoxypsoralen), a psoralen, inhibits cell replication.
Targets(IC50)	Autophagy,Cytochromes P450
In vitro	In C57BL/6J mice induced with acetaminophen (APAP), Bergapten exhibits antioxidant activity and mitigates hepatotoxicity.
In vivo	In both tamoxifen-sensitive and -resistant breast cancer cells, Bergapten eliminates ERα and reprograms membrane signaling targeted at ERα, thereby enhancing the potential for cell mitosis.
Cell Research	5-Methoxypsoralen (5-MOP) is dissolved in DMSO and stored, and then diluted with appropriate medium (final DMSO <0.1%) before use[1]. The human colon adenocarcinoma cell line (COLO 205, from a 70-year-old male Caucasian) are placed into 75-cm <sup>2</sup> tissue culture flasks and grown in RPMI1640 medium, supplemented with 10% fetal bovine serum, containing penicillin and streptomycin (100 Ig/mL) and 1 mM glutamine, at 37°C in a humidified atmosphere of 5% CO <sub>2</sub> and 95% O <sub>2</sub> . The human stomach adenocarcinoma cell line are placed into 75-cm <sup>2</sup> tissue culture flasks and grown in RPMI 1640 medium, supplemented with 10% fetal bovine serum, containing penicillin and streptomycin (100 Ig/mL) and 1 mM glutamine, at 37°C in a humidified atmosphere of 5% CO <sub>2</sub> and 95% O <sub>2</sub> . SC-M1 and COLO 205 cells are treated with different concentrations of 5-MOP (0.05, 0.5, 5, 10, 25 and 50 mM) and incubated for 72 h for the dose-effect study of 5-MOP on NAT activity. To determine the time-course effect of 0.5 mM 5-MOP on NAT activity, the cells are incubated at 37AC and harvested at 12, 24, 48 and 72 h, respectively. 5-MOP is dissolved in DMSO and the final concentration of vehicle is <0.1%. Only DMSO (solvent) is added for the control regimen[1].

## Solubility Information

Solubility	Ethanol: < 1 mg/mL (insoluble or slightly soluble), H <sub>2</sub> O: < 1 mg/mL (insoluble or slightly soluble), DMSO: 20 mg/mL (92.51 mM),Sonication is recommended. (< 1 mg/ml refers to the product slightly soluble or insoluble)
------------	--

### Preparing Stock Solutions

	1mg	5mg	10mg
1 mM	4.6256 mL	23.1278 mL	46.2556 mL
5 mM	0.9251 mL	4.6256 mL	9.2511 mL
10 mM	0.4626 mL	2.3128 mL	4.6256 mL
50 mM	0.0925 mL	0.4626 mL	0.9251 mL

Please select the appropriate solvent to prepare the stock solution, according to the solubility of the product in different solvents. Please use it as soon as possible.

### Reference

Panno ML, et al. Breast Cancer Res Treat, 2012, 136(2), 443-455.

Gao S, Zou X, Wang Z, et al. Bergapten attenuates microglia-mediated neuroinflammation and ischemic brain injury by targeting Kv1. 3 and Carbonyl reductase 1. European Journal of Pharmacology. 2022: 175242.

Liu WX, et al. World J Gastroenterol, 2012, 18(18), 2197-2202.

**Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins**

**This product is for Research Use Only. Not for Human or Veterinary or Therapeutic Use**

Tel:781-999-4286    E\_mail:info@targetmol.com    Address:36 Washington Street,Wellesley Hills,MA 02481