


Tacrolimus

Chemical Properties

CAS No. :	104987-11-3
Formula:	C ₄₄ H ₆₉ NO ₁₂
Molecular Weight:	804.02
Appearance:	no data available
Storage:	Powder: -20°C for 3 years In solvent: -80°C for 1 year



Biological Description

Description	Tacrolimus (Fujimycin) can bind FKBP12 to form a high-affinity complex (K _i : 0.2 nM) which inhibits the activity of the calcium/calmodulin-dependent protein phosphatase.
Targets(IC50)	Antibacterial,Antibiotic,Autophagy,mTOR,Phosphatase
In vitro	FKBP12 was first isolated from the cytosol of Jurkat T cells, as an abundant, high-affinity receptor for Tacrolimus (FK506; K _d : 0.2 nM) [1]. FK506 (100?1,000 µg/l) significantly promoted the proliferation of MH3924A cells. The invasiveness of MH3924A cells was significantly enhanced following treatment with FK506 [2]. FK506 specifically inhibits cellular calcineurin at drug concentrations that inhibit interleukin 2 productions in activated T cells [3].
In vivo	The liver of rats of the normal saline (NS) group was large, with an average weight at 15.56±11.17 g, and the liver of rats of the FK506 group was oversize, with an average weight at 28.19±3.89 g. The rate of lymph node metastasis, as well as the number of pulmonary nodules, were significantly increased in the FK506 compared to the NS group [2]. Behavioral pain assessment revealed an increase in the paw and tail withdrawal threshold in tacrolimus-treated groups against hyperalgesic and allodynic stimuli as compared to the sham control group [4].
Cell Research	Tumor cell proliferation was determined by the MTT. Briefly, after MH3924A cells had reached the logarithmic growth phase, a 0.2-ml cell suspension at 1×10 ⁴ cells/ml was added into each well of a 96-well plate and cultured in DMEM with 10% FBS, 10 µg/l vascular endothelial growth factor and 0.1 g/l heparin for 24 h. When adherent growth was established, different concentrations of FK506 (10, 100 and 1,000 µg/l), AMD3100 (10, 50 and 100 µg/l) and FK506 (0 and 100 µg/l) + AMD3100 (0, 10, 50 and 100 µg/l) were added into the plates. Untreated cells cultured in medium alone were used as controls. After culturing for 48 h, 10 µl MTT (5 g/l) was added, and each well was incubated for 6 h; next, 150 µl/well dimethyl sulfoxide was added, followed by measurements of the absorbance at 570 nm on a spectrophotometer reader. Each well was measured three times, and each sample was assayed in triplicate [2].
Animal Research	Experiments were performed in 16 healthy August Copenhagen Irish rats (male, 16–20 weeks, weighing 240–300 g). The rat model of liver tumor was established as follows: First, MH3924A cells were collected and injected into the alar skin of rats. The tumors were removed from alar skin when grown to 2×1×1 mm ³ , and intrahepatic tumor implantation of rats was performed under aseptic conditions as described previously

(25,26). Five days later, rats were randomly divided into two groups: one group was subcutaneously injected with normal saline for 14 days (NS group, n=8, 3 mg/kg/day), and the second group was subcutaneously injected with FK506 for 14 days (FK506 group, n=8, 0.3 mg/kg/day). Forty days following implantation, rats were sacrificed, and the weight of tumor, the volume of the fluid in the ascites, the incidence of lymphatic metastasis in the abdominal cavity and of abdominal wall metastasis were measured. In addition, the lungs were irrigated with 15% Indian ink, followed by counting of the number of metastatic nodules in the lung. The tumor and adjacent tissues, as well as healthy liver tissues, were harvested and preserved in 4% formalin for later use [2].

Solubility Information

Solubility	Ethanol: 80.4 mg/mL (100 mM), Sonication is recommended. DMSO: 45 mg/mL (55.97 mM), Sonication is recommended. 10% DMSO+40% PEG300+5% Tween 80+45% Saline: 8.04 mg/mL (10 mM), Solution. (< 1 mg/mL refers to the product slightly soluble or insoluble)
------------	--

Preparing Stock Solutions

	1mg	5mg	10mg
1 mM	1.2438 mL	6.2188 mL	12.4375 mL
5 mM	0.2488 mL	1.2438 mL	2.4875 mL
10 mM	0.1244 mL	0.6219 mL	1.2438 mL
50 mM	0.0249 mL	0.1244 mL	0.2488 mL

Please select the appropriate solvent to prepare the stock solution, according to the solubility of the product in different solvents. Please use it as soon as possible.

Reference

- Zhu H, et al. Tacrolimus promotes hepatocellular carcinoma and enhances CXCR4/SDF-1 α expression in vivo. *Mol Med Rep.* 2014 Aug;10(2):585-92.
- Wang W, Ren S, Lu Y, et al. Inhibition of Syk promotes chemical reprogramming of fibroblasts via metabolic rewiring and H2S production. *The EMBO Journal.* 2021 Jun 1;40(11):e106771. doi: 10.15252/embj.2020106771. Epub 2021 Apr 28.
- Chang Z, Zhang Q, Hu Q, et al. Tannins in Terminalia bellirica inhibits hepatocellular carcinoma growth via re-educating tumor-associated macrophages and restoring CD8 $^{+}$ T cell function. *Biomedicine & Pharmacotherapy.* 2022, 154: 113543.
- Bennett J, et al. Tacrolimus Modulates TGF- β Signaling to Induce Epithelial-Mesenchymal Transition in Human Renal Proximal Tubule Epithelial Cells. *J Clin Med.* 2016 Apr 26;5(5):50.
- Manojlovic Z, et al. Tacrolimus (FK506) prevents early stages of ethanol induced hepatic fibrosis by targeting LARP6 dependent mechanism of collagen synthesis. *PLoS One.* 2013 Jun 3;8(6):e65897.
- Huang, Jing, et al. Osteopontin isoform c promotes the survival of cisplatin-treated NSCLC cells involving NFATc2-mediated suppression on calcium-induced ROS levels.. *BMC cancer.* 21.1 (2021): 1-14.
- Muthuraman A, Sood S. Pharmacological evaluation of tacrolimus (FK-506) on ischemia reperfusion induced vasculatic neuropathic pain in rats. *J Brachial Plex Peripher Nerve Inj.* 2010 Jun 7;5:13.
- Yuwei He, et al. Drug targeting through platelet membrane-coated nanoparticles for the treatment of rheumatoid arthritis. *Nano Res.* 30 June 2018.
- Wang W, Ren S, Lu Y, et al. Inhibition of Syk promotes chemical reprogramming of fibroblasts via metabolic rewiring and H2S production. *The EMBO Journal.* 2021: e106771.

Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins

This product is for Research Use Only· Not for Human or Veterinary or Therapeutic Use

Tel:781-999-4286 E_mail:info@targetmol.com Address:36 Washington Street,Wellesley Hills,MA 02481