


BX471

## Chemical Properties

CAS No. : 217645-70-0  
 Formula: C<sub>21</sub>H<sub>24</sub>ClFN<sub>4</sub>O<sub>3</sub>  
 Molecular Weight: 434.89  
 Appearance: no data available  
 Storage: Powder: -20°C for 3 years | In solvent: -80°C for 1 year



## Biological Description

Description	BX471 (BX 471) is a potent, selective non-peptide CCR1 antagonist.
Targets(IC50)	CCR
In vitro	BX 471 is a potent functional antagonist based on its ability to inhibit a number of CCR1-mediated effects including Ca <sup>2+</sup> mobilization, increase in extracellular acidification rate, CD11b expression, and leukocyte migration. BX 471 demonstrates a greater than 10,000-fold selectivity for CCR1 compared with 28 G-protein-coupled receptors[1]. BX471 is also able to displace 125I-MIP-1 $\alpha$ /CCL3 binding to mouse CCR1 in a concentration-dependent manner with a K <sub>i</sub> of 215 $\pm$ 46 nM. Increasing concentrations of BX471 inhibits the Ca <sup>2+</sup> transients induced by MIP-1 $\alpha$ /CCL3 in both human and mouse CCR1 with IC <sub>50</sub> of 5.8 $\pm$ 1 nM and 198 $\pm$ 7 nM, respectively[2]. BX471 (0.1-10 $\mu$ M) shows a dose-dependent inhibition of RANTES-mediated and shear-resistant adhesion on IL-1 $\beta$ -activated microvascular endothelium in shear flow in isolated blood monocytes. BX471 also inhibits the RANTES-mediated adhesion of T lymphocytes to activated endothelium[4].
In vivo	BX 471, administered orally (p.o.) or intravenously (i.v.) at a dosage of 4 mg/kg, demonstrates an oral bioavailability of 60% in dogs and effectively mitigates symptoms in a rat model of multiple sclerosis, specifically the experimental allergic encephalomyelitis model[1]. When given subcutaneously (s.c.) at 20 mg/kg, BX 471 achieves peak plasma concentrations of 9 $\mu$ M within approximately 30 minutes, subsequently decreasing to about 0.4 $\mu$ M after 2 hours and further dropping to 0.1 $\mu$ M or below from 4 to 8 hours. Treatment with 20 mg/kg BX 471 for 10 days in mice results in a 55% reduction in interstitial CD45 positive leukocytes. Although the effect on CCR5-positive CD8 cells in peripheral blood is marginally significant, BX 471 notably reduces FSP1-positive cells in UUO (unilateral ureteral obstruction) kidney conditions by 65% compared to controls[2], and pre-treatment with BX 471 significantly diminishes macrophage and neutrophil accumulation in the kidney following ischemia-reperfusion injury[3].
Kinase Assay	Kinase activity assays: In vitro activities of purified GST-NUAK1 and GST-NUAK1[A195T] are measured using Cerenkov counting of incorporation of radioactive <sup>32</sup> P from [ $\gamma$ - <sup>32</sup> P] ATP into Sakamototide substrate peptide. Reactions are carried out in a 50 $\mu$ L reaction volume for 30 min at 30°C and reactions are terminated by spotting 40 $\mu$ L of the reaction mix on to P81 paper and immediately immersing in 50 mM orthophosphoric acid. Samples are washed three times in 50 mM orthophosphoric acid followed by a single

acetone rinse and air drying. The kinase-mediated incorporation of [ $\gamma$ - $^{32}$ P]ATP into Sakamototide is quantified by Cerenkov counting. One unit of activity is defined as that which catalyses the incorporation of 1 nmol of [ $^{32}$ P]phosphate into the substrate over 1 h.

## Cell Research

Briefly, dermal microvascular endothelial cells grown to confluence in Petri dishes are stimulated with IL-1 $\beta$  (10 ng/mL) for 12 h followed by pre-incubation with RANTES (10 nM) for 30 min at 37°C just prior to assay. The plates are assembled as the lower wall in a parallel wall flow chamber and mounted on the stage of an Olympus IMT-2 inverted microscope with  $\times 20$  and  $\times 40$  phase-contrast objectives. Isolated human blood monocytes are isolated and resuspended at  $5 \times 10^5$  cells/mL in assay buffer (HBSS) containing 10 mM HEPES, pH 7.4 and 0.5% human serum albumin. Shortly before the assay, 1 mM Mg $^{2+}$  and 1 mM Ca $^{2+}$  are added. The cell suspensions are kept in a heating block at 37°C during the assay and perfused into the flow chamber at a rate of 1.5 dyn/cm $^2$  for 5 min. For inhibition experiments, monocytes are preincubated with BX471 at different concentrations (0.1–10  $\mu$ M) or a Me2SO control for 10 min at 37°C. The number of firmly adherent cells after 5 min is quantitated in multiple fields (at least five per experiment) by analysis of images recorded with a long integration JVC 3CCD video camera and a JVC SR L 900 E video recorder and are expressed as cells/mm $^2$ . The type of adhesion analyzed is restricted to primary, i.e. direct interactions of monocytes with endothelium.

## Solubility Information

## Solubility

DMSO: 100 mg/mL (229.94 mM), Sonication is recommended.  
( $< 1$  mg/mL refers to the product slightly soluble or insoluble)

## Preparing Stock Solutions

	1mg	5mg	10mg
1 mM	2.2994 mL	11.4972 mL	22.9943 mL
5 mM	0.4599 mL	2.2994 mL	4.5989 mL
10 mM	0.2299 mL	1.1497 mL	2.2994 mL
50 mM	0.046 mL	0.2299 mL	0.4599 mL

Please select the appropriate solvent to prepare the stock solution, according to the solubility of the product in different solvents. Please use it as soon as possible.

### Reference

- Liang M, et al. Identification and characterization of a potent, selective, and orally active antagonist of the CC chemokine receptor-1. J Biol Chem. 2000 Jun 23;275(25):192000-8.
- Yang J Y, Zhang J, Lu R, et al. T cell-derived exosomes induced macrophage inflammatory protein-1 $\alpha$ / $\beta$  drive the trafficking of CD8+ T cells in oral lichen planus. Journal of Cellular and Molecular Medicine. 2020
- Anders HJ, et al. A chemokine receptor CCR-1 antagonist reduces renal fibrosis after unilateral ureter ligation. J Clin Invest. 2002 Jan;109(2):251-9.
- Furuichi K, et al. Chemokine receptor CCR1 regulates inflammatory cell infiltration after renal ischemia-reperfusion injury. J Immunol. 2008 Dec 15;181(12):8670-6.
- Horuk R, et al. A non-peptide functional antagonist of the CCR1 chemokine receptor is effective in rat heart transplant rejection. J Biol Chem. 2001 Feb 9;276(6):4199-204.
- Yang J Y, Zhang J, Lu R, et al. T cell-derived exosomes induced macrophage inflammatory protein-1 $\alpha$ / $\beta$  drive the trafficking of CD8+ T cells in oral lichen planus[J]. Journal of cellular and molecular medicine. 2020

**Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins**

**This product is for Research Use Only· Not for Human or Veterinary or Therapeutic Use**

Tel:781-999-4286    E\_mail:info@targetmol.com    Address:36 Washington Street,Wellesley Hills,MA 02481