


Nocodazole

Chemical Properties

CAS No. :	31430-18-9
Formula:	C ₁₄ H ₁₁ N ₃ O ₃ S
Molecular Weight:	301.32
Appearance:	no data available
Storage:	Powder: -20°C for 3 years In solvent: -80°C for 1 year



Biological Description

Description	Nocodazole (Oncodazole) is a reversible inhibitor of microtubule polymerization and an inhibitor of Bcr-Abl. Nocodazole has antitumor activity, blocks the cell cycle and induces apoptosis.
Targets(IC50)	Apoptosis, Microtubule Associated, Bcr-Abl, Autophagy, CRISPR/Cas9
In vitro	<p>METHODS: Erythrocytes were treated with Nocodazole (15-60 µg/mL) for 48 h. Phosphatidylserine was detected using Annexin-V-FITC.</p> <p>RESULTS: Nocodazole treatment increased the percentage of phosphatidylserine exposed to erythrocytes, reaching statistical significance at 30 µg/mL. [1]</p> <p>METHODS: hESCs cells were treated with Nocodazole (100 ng/mL) for 16 h and cell cycle profiles were analyzed using Flow Cytometry.</p> <p>RESULTS: Nocodazole treatment synchronized the cell cycle (>90% of cells in G2/M), while cells remained synchronized after release and moved evenly through the cell cycle for 24 h. At 2 h after removal of Nocodazole, the cells entered the G1 phase, with 70% of the cells in the G1 phase after 4 h, and 80% in the S phase after 12 h. The cells were treated with Nocodazole (100 ng/mL) for 16 h, and the cell cycle was analyzed by Flow Cytometry. [2]</p>
In vivo	<p>METHODS: To detect anti-tumor activity in vivo, Nocodazole (12 mg/kg three times a week) and dexamethasone (2 mg/kg twice a week) were injected intraperitoneally for fifteen days into immunodeficient mice bearing myeloma H929.</p> <p>RESULTS: Nocodazole in combination with dexamethasone significantly inhibited myeloma tumor growth and prolonged survival. [3]</p> <p>METHODS: To study the effects on intestinal mucositis, Nocodazole (5 mg/kg) and ketoconazole (50 mg/kg) were administered intraperitoneally three times a week for six weeks to nude mice harboring the human colorectal cancer tumor COLO 205.</p> <p>RESULTS: The antitumor effect of Nocodazole was significantly enhanced after six weeks of ketoconazole treatment. [4]</p>
Cell Research	Nocodazole is dissolved in a final concentration of 0.05% DMSO. Proteins are loaded at 50 µg/lane and separated by 12% (w:v) sodium dodecyl sulfate-polyacrylamide gel electrophoresis, blotted, and probed with antibodies for cyclin E, p53, p21/CIP1, p27/KIP1, glyceraldehyde 3-phosphate dehydrogenase (GAPDH), cyclin A, cyclin D1, cyclin D3, cyclin B, CDK2, CDK4, and cytochrome C. Immunoreactive bands are visualized by incubating with the colorigenic substrates nitroblue tetrazolium and 5-bromo-4-chloro-3-indolyl-phosphate. The expression of GAPDH is used as the control for equal

protein loading.

Solubility Information

Solubility	DMSO: 12.5 mg/mL (41.48 mM),Sonication is recommended. (< 1 mg/ml refers to the product slightly soluble or insoluble)
------------	---------------------------------------------------------------------------------------------------------------------------

Preparing Stock Solutions

	1mg	5mg	10mg
1 mM	3.3187 mL	16.5937 mL	33.1873 mL
5 mM	0.6637 mL	3.3187 mL	6.6375 mL
10 mM	0.3319 mL	1.6594 mL	3.3187 mL
50 mM	0.0664 mL	0.3319 mL	0.6637 mL

Please select the appropriate solvent to prepare the stock solution, according to the solubility of the product in different solvents. Please use it as soon as possible.

Reference

- Signoretto E, et al. Nocodazole Induced Suicidal Death of Human Erythrocytes. *Cell Physiol Biochem*. 2016;38(1): 379-92.
- Bai Y, Wang W, Li S, et al. C1QBP Promotes Homologous Recombination by Stabilizing MRE11 and Controlling the Assembly and Activation of MRE11/RAD50/NBS1 Complex. *Molecular Cell*. 2019
- Bai Y, Wang W, Li S, et al. C1QBP Promotes Homologous Recombination by Stabilizing MRE11 and Controlling the Assembly and Activation of MRE11/RAD50/NBS1 Complex[J]. *Molecular Cell*. 2019.
- Yiangou L, et al. Method to Synchronize Cell Cycle of Human Pluripotent Stem Cells without Affecting Their Fundamental Characteristics. *Stem Cell Reports*. 2019 Jan 8;12(1):165-179.
- Moreira G G, Cantrelle F X, Quezada A, et al. Dynamic interactions and Ca²⁺-binding modulate the holdase-type chaperone activity of S100B preventing tau aggregation and seeding. *Nature Communications*. 2021, 12(1): 1-16
- Reyes G X, Kolodziejczak A, Devakumar L J P S, et al. Ligation of newly replicated DNA controls the timing of DNA mismatch repair. *Current Biology*.. 2021 Mar 22;31(6):1268-1276.e6. doi: 10.1016/j.cub.2020.12.018. Epub 2021 Jan 7.
- Feng R, et al. Targeting the microtubular network as a new antimyeloma strategy. *Mol Cancer Ther*. 2011 Oct;10 (10):1886-96.
- Wang YJ, et al. Ketoconazole potentiates the antitumor effects of nocodazole: In vivo therapy for human tumor xenografts in nude mice. *Mol Carcinog*. 2002 Aug;34(4):199-210.
- Li X, Tang H, Huang X, et al. Rigidity-Dependent Placental Cells Uptake of Silk-Based Microcapsules. *Macromolecular bioscience*. 2019: 1900105.
- Cui L, Xu Q, Zhang P, et al. Biochemical reprogramming of active-targeting and redox responsive nanoparticles for enhancing selective internalization. *Materials Letters*. 2023: 134379.
- Wang YJ, et al. *Mol Carcinog*. 2002, 34(4), 199-210.
- Wang J, Chen Y, Li S, et al. PP2A inhibition causes synthetic lethality in BRCA2-mutated prostate cancer models via reactivating spindle assembly checkpoint. *The Journal of Clinical Investigation*. 2023
- Signoretto E, et al. Nocodazole Induced Suicidal Death of Human Erythrocytes. *Cell Physiol Biochem*. 2016;38(1): 379-92.
- Zhang JP, et al. Efficient precise knockin with a double cut HDR donor after CRISPR/Cas9-mediated double-stranded DNA cleavage. *Genome Biol*. 2017 Feb 20;18(1):35.
- Li X, Tang H, Huang X, et al. Rigidity-Dependent Placental Cells Uptake of Silk-Based Microcapsules[J]. *Macromolecular bioscience*. 2019: 1900105.
- Reyes G X, Kolodziejczak A, Devakumar L J P S, et al. Ligation of newly replicated DNA controls the timing of DNA mismatch repair[J]. *Current Biology*.

Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins

This product is for Research Use Only. Not for Human or Veterinary or Therapeutic Use

Tel: 781-999-4286 E_mail: info@targetmol.com Address: 36 Washington Street, Wellesley Hills, MA 02481