


Puerarin

Chemical Properties

CAS No. :	3681-99-0
Formula:	C ₂₁ H ₂₀ O ₉
Molecular Weight:	416.38
Appearance:	no data available
Storage:	Powder: -20°C for 3 years In solvent: -80°C for 1 year



Biological Description

Description	Puerarin (Kakonein), also known as Kakonein, is a member of the class of compounds known as isoflavonoid C-glycosides. It is a 5-HT _{2C} receptor antagonist.
Targets(IC ₅₀)	5-HT Receptor
In vitro	Administering 300 mg/kg of Puerarin daily through oral ingestion to rats on a high-cholesterol diet significantly reduces the increase in total cholesterol levels in both serum and liver caused by the high-cholesterol diet.
In vivo	Puerarin, at a dose of 25 μ M, induces dose-dependent growth inhibition in HT-29 cells. This effect is accompanied by an increase in bax and a decrease in c-myc and bcl-2.
Kinase Assay	STAT3-dependent dual-luciferase assay: HCT-116 cells are transiently transfected with reporter plasmid having the STAT3-binding element for regulating luciferase assay. Cells are treated with Cryptotanshinone for 24 hours at a concentration range of 0.2 to 50 μ M. After treatment, cells are harvested in 20 μ L of passive lysis buffer and luciferase activity is evaluated by the Dual Luciferase Reporter Assay kit on Wallac Victor2. The concentration of Cryptotanshinone that inhibits the luciferase activity by 50% represents IC ₅₀ value.
Cell Research	RAW264.7 cells are maintained at subconfluence in 95% air and 5% CO ₂ humidified atmosphere maintained at 37°C. The medium used for routine subculture is Dulbecco's Modified Eagle's Medium supplemented with 10% fetal bovine serum, penicillin (100 units/mL) and streptomycin (100 μ g/mL). An MTT assay is used to measure the viability of the cells after treatment with puerarin. After the supernatants are removed for nitrite determination, cells are incubated at 37°C with MTT (0.05 mg/mL) for 4 h, and the optical density is measured at 540 nm. The concentrations of puerarin are 10, 20, 40 and 100 μ M[1].

Solubility Information

Solubility	DMSO: 50 mg/mL (120.08 mM), Sonication is recommended. Ethanol: < 1 mg/mL (insoluble or slightly soluble), H ₂ O: < 1 mg/mL (insoluble or slightly soluble), (< 1 mg/mL refers to the product slightly soluble or insoluble)
------------	--

Preparing Stock Solutions

	1mg	5mg	10mg
1 mM	2.4017 mL	12.0083 mL	24.0165 mL
5 mM	0.4803 mL	2.4017 mL	4.8033 mL
10 mM	0.2402 mL	1.2008 mL	2.4017 mL
50 mM	0.048 mL	0.2402 mL	0.4803 mL

Please select the appropriate solvent to prepare the stock solution, according to the solubility of the product in different solvents. Please use it as soon as possible.

Reference

Overstreet DH, et al. Pharmacol Biochem Behav, 2003, 75(3), 619-625.

Wang X, He K, Ma L, et al. Puerarin attenuates isoproterenol-induced myocardial hypertrophy via inhibition of the Wnt/ β -catenin signaling pathway. Molecular Medicine Reports. 2022, 26(4): 1-13.

Liang Y, Xu Z, Wu X, et al. Inhibition of hyperpolarization-activated cyclic nucleotide-gated channels with natural flavonoid quercetin. Biochemical and Biophysical Research Communications. 2020

Yan LP, et al. Life Sci, 2006, 79(4), 324-330.

Yu Z, et al. Cancer Lett, 2006, 238(1), 53-60.

Wang K, Sun Y, Zhu K, et al. Anti-Pyroptosis Biomimetic Nanoplatfrom loading Puerarin for Myocardial Infarction Repair: From Drug Discovery to Drug Delivery. Biomaterials. 2024: 122890.

Xu Y, Liang H, Mao X, et al. Puerarin alleviates apoptosis and inflammation in kidney stone cells via the PI3K/AKT pathway: Network pharmacology and experimental verification. Journal of Cellular and Molecular Medicine. 2024, 28(20): e70180.

Liang Y, Xu Z, Wu X, et al. Inhibition of hyperpolarization-activated cyclic nucleotide-gated channels with natural flavonoid quercetin[J]. Biochemical and Biophysical Research Communications. 2020

Xu Y, Cai Q, Zhao C, et al. Gegen Qinlian Decoction Attenuates Colitis-Associated Colorectal Cancer via Suppressing TLR4 Signaling Pathway Based on Network Pharmacology and In Vivo/In Vitro Experimental Validation. Pharmaceuticals. 2024, 18(1): 12.

Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins

This product is for Research Use Only · Not for Human or Veterinary or Therapeutic Use

Tel: 781-999-4286 E_mail: info@targetmol.com Address: 36 Washington Street, Wellesley Hills, MA 02481