


Osthole

Chemical Properties

CAS No. :	484-12-8
Formula:	C ₁₅ H ₁₆ O ₃
Molecular Weight:	244.29
Appearance:	no data available
Storage:	Powder: -20°C for 3 years In solvent: -80°C for 1 year



Biological Description

Description	Osthole (Osthonolide), a potential inhibitor of histamine H1 receptor, has been shown to stimulate osteoblast proliferation and differentiation.
Targets(IC50)	Apoptosis,Parasite,Histamine Receptor,HBV
In vitro	The LD50 values of Osthole in various animals have been determined as follows: in rats, it exceeds 125 mg/kg when administered intravenously (i.v.); in rabbits, it is over 35 mg/kg (i.v.); and in dogs, it surpasses 150 mg/kg (i.v.).
In vivo	Osthonolide, when introduced to pre-washed rabbit platelets, inhibits platelet aggregation and ATP release induced by ADP, arachidonic acid, platelet-activating factor (PAF), collagen, ionophore A23187, and thrombin. In human osteoblasts, osthonolide promotes cell differentiation through the bone morphogenetic protein-2 / p38 and extracellular signal-regulated kinase 1/2 pathways. Additionally, in cell cultures, osthonolide suppresses the secretion of the hepatitis B virus by enhancing the glycosylation of HBsAg.
Cell Research	Osthole is dissolved in 96% ethyl alcohol and then sterilized through a 0.22 µg/mL filter and stored at 4°C as stock solutions for later dilution[1]. Peripheral blood samples are collected from participants between 7.00 and 9.00 a.m. on the first study day and these are concentrated in grouping tubes with K3EDTA. Fresh PBMCs are then prepared. Isolated cells are seeded on 24-well plates at 1×10 ⁶ per well with RPMI-1640 and supplemented with 1% heat inactivated human AB serum, 1% gentamicin and 0.25% PHA. Active reagents are added to each well after 24 h and pure medium formed the control for each substance. Cells are then harvested after a further three days[1].

Solubility Information

Solubility	Ethanol: 46 mg/mL (188.3 mM),Sonication is recommended. DMSO: 55 mg/mL (225.14 mM),Sonication is recommended. H2O: < 1 mg/mL (insoluble or slightly soluble), (< 1 mg/ml refers to the product slightly soluble or insoluble)
------------	--

Preparing Stock Solutions

	1mg	5mg	10mg
1 mM	4.0935 mL	20.4675 mL	40.935 mL
5 mM	0.8187 mL	4.0935 mL	8.187 mL
10 mM	0.4093 mL	2.0467 mL	4.0935 mL
50 mM	0.0819 mL	0.4093 mL	0.8187 mL

Please select the appropriate solvent to prepare the stock solution, according to the solubility of the product in different solvents. Please use it as soon as possible.

Reference

Ko FN, et al. Thromb Haemost, 1989, 62(3), 1996-1999.

Geng W, Guo X, Zhang L, et al. Resveratrol inhibits proliferation, migration and invasion of multiple myeloma cells via NEAT1-mediated Wnt/ β -catenin signaling pathway Resveratrol inhibits proliferation, migration and invasion of multiple myeloma cells via NEAT1-mediated Wnt/ β -catenin signaling pathway. Biomedicine & Pharmacotherapy. 2018 Nov;107:484-494.

Qu Y, Sun X, Wei N, et al. Inhibition of cutaneous heat-sensitive Ca^{2+} -permeable transient receptor potential vanilloid 3 channels alleviates UVB-induced skin lesions in mice. The FASEB Journal. 2023, 37(12): e23309.

Kuo PL, et al. J Pharmacol Exp Ther, 2005, 314(3), 1290-1299.

Xu Y, Qu Y, Zhang C, et al. Selective inhibition of overactive warmth-sensitive Ca^{2+} -permeable TRPV3 channels by antispasmodic agent flopropione for alleviation of skin inflammation. Journal of Biological Chemistry. 2023: 105595.

Huang RL, et al. Hepatology, 1996, 24(3), 508-515.

Geng W, Guo X, Zhang L, et al. Resveratrol inhibits proliferation, migration and invasion of multiple myeloma cells via NEAT1-mediated Wnt/ β -catenin signaling pathway[J]. Biomedicine & Pharmacotherapy. 2018 Nov;107:484-494.

Li S H, Li M Y, Yuan T T, et al. Osthole Activates the Cholinergic Anti-Inflammatory Pathway via $\alpha 7\text{nAChR}$ Upregulation to Alleviate Inflammatory Responses. Chemistry & Biodiversity. 2024: e202400290.

Yu X, Jiang J, Li C, et al. Osthole ameliorates wear particle-induced osteogenic impairment by mitigating endoplasmic reticulum stress via PERK signaling cascade. Molecular Medicine. 2024, 30: 266.

Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins

This product is for Research Use Only · Not for Human or Veterinary or Therapeutic Use

Tel: 781-999-4286 E_mail: info@targetmol.com Address: 36 Washington Street, Wellesley Hills, MA 02481