


RG13022

Chemical Properties

CAS No. : 136831-48-6
Formula: C₁₆H₁₄N₂O₂
Molecular Weight: 266.29
Appearance: no data available
Storage: Powder: -20°C for 3 years | In solvent: -80°C for 1 year



Biological Description

Description	RG13022 (Tyrphostin RG13022) is a tyrosine kinase inhibitor; inhibits the autophosphorylation reaction of the EGF receptor (IC ₅₀ : 4 μM).
Targets(IC ₅₀)	EGFR
In vitro	RG13022 suppresses EGF-stimulated cancer cell proliferation and inhibits EGF receptor autophosphorylation in immunoprecipitates (IC ₅₀ : 4 μM) in a cell-free reaction. It inhibits colony formation (IC ₅₀ : 1 μM) and DNA synthesis (IC ₅₀ : 3 μM) in HER 14 cells stimulated by 50 ng/mL EGF, dose-dependently. Additionally, RG13022 inhibits growth stimulated by insulin, insulin-like growth factor I, insulin-like growth factor II, or transforming growth factor alpha. It completely blocks estrogen-stimulated phosphorylation of the EGF receptor and estrogen-induced cell proliferation, indicating that functioning TK pathways are essential for estrogen action.
In vivo	RG13022 suppresses tumor growth and extends the lifespan of tumor-bearing nude mice.
Cell Research	40 mM stock solutions of RG13022 is made in 100% DMSO and diluted with the culture medium before addition to the cells. MH-85 cells and HER 14 cells are plated in culture medium in the presence or absence of increasing concentrations of RG-13022 or RG-14620 for 10 days. At the end of culture, the cells are fixed with 4% (v/v) formaldehyde and stained with hematoxylin. Numbers of colonies including more than 20 cells in each well are counted under the microscope[1].

Solubility Information

Solubility	DMSO: 50 mg/mL (187.77 mM),Sonication is recommended. (< 1 mg/ml refers to the product slightly soluble or insoluble)
------------	--

Preparing Stock Solutions

	1mg	5mg	10mg
1 mM	3.7553 mL	18.7765 mL	37.553 mL
5 mM	0.7511 mL	3.7553 mL	7.5106 mL
10 mM	0.3755 mL	1.8777 mL	3.7553 mL
50 mM	0.0751 mL	0.3755 mL	0.7511 mL

Please select the appropriate solvent to prepare the stock solution, according to the solubility of the product in different solvents. Please use it as soon as possible.

Reference

Yoneda T, et al. The antiproliferative effects of tyrosine kinase inhibitors tyrphostins on a human squamous cell carcinoma in vitro and in nude mice. *Cancer Res.* 1991 Aug 15;51(16):4430-5.

Nguyen D J M, Theodoropoulos G, Li Y Y, et al. Targeting the kynurenine pathway for the treatment of cisplatin resistant lung cancer. *Molecular Cancer Research.* 2019: molcanres. 0239.2019

Reddy KB, et al. Inhibition of breast cancer cell growth in vitro by a tyrosine kinase inhibitor. *Cancer Res.* 1992 Jul 1; 52(13):3636-41.

Nguyen D J M, Theodoropoulos G, Li Y Y, et al. Targeting the kynurenine pathway for the treatment of cisplatin-resistant lung cancer[J]. *Molecular Cancer Research.* 2020, 18(1): 105-117.

Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins

This product is for Research Use Only · Not for Human or Veterinary or Therapeutic Use

Tel:781-999-4286 E_mail:info@targetmol.com Address:36 Washington Street,Wellesley Hills,MA 02481