

2-heptyl-3-hydroxy-4(1H)-Quinolone

Chemical Properties

CAS No. : 108985-27-9


Formula: C16H21NO2

Molecular Weight: 259.34

Appearance: Solid

Storage: store at low temperature, store under nitrogen

Powder: -20°C for 3 years | In solvent: -80°C for 1 year



Biological Description

Description	2-heptyl-3-hydroxy-4(1H)-Quinolone (Pseudomonas Quinolone Signal) is a quorum-sensing signalling molecule produced by <i>Pseudomonas aeruginosa</i> in response to increased cell density. It increases the expression of the <i>lasB</i> gene in <i>P. aeruginosa</i> , the secretion of the metabolites pyocyanin and the lectin PA-IL, as well as increasing biofilm production in <i>P. aeruginosa</i> populations. It also reduces the iron content of the <i>P. aeruginosa</i> growth medium when used at a concentration of 40 µM and acts as an iron chelator in iron sulphate solutions.
Targets(IC50)	Others, Antibacterial
In vitro	2-heptyl-3-hydroxy-4(1H)-Quinolone (60 µM) increases secretion of the metabolite pyocyanin and the lectin PA-IL, as well as increases biofilm production in <i>P. aeruginosa</i> populations.[2] 2-heptyl-3-hydroxy-4(1H)-Quinolone (40 µM) reduces iron levels in <i>P. aeruginosa</i> growth media when used at a concentration and acts as an iron chelator in a Fe(III)-sulfate solution.[3]

Solubility Information

Solubility	10% DMSO+40% PEG300+5% Tween 80+45% Saline: 2 mg/mL (7.71 mM), Sonication is recommended. Methanol: Soluble DMSO: 12 mg/mL (46.27 mM), Sonication is recommended. (< 1 mg/ml refers to the product slightly soluble or insoluble)
------------	--

Preparing Stock Solutions

	1mg	5mg	10mg
1 mM	3.8559 mL	19.2797 mL	38.5594 mL
5 mM	0.7712 mL	3.8559 mL	7.7119 mL
10 mM	0.3856 mL	1.928 mL	3.8559 mL
50 mM	0.0771 mL	0.3856 mL	0.7712 mL

Please select the appropriate solvent to prepare the stock solution, according to the solubility of the product in different solvents. Please use it as soon as possible.

Reference

Pesci EC, et al. Quinolone signaling in the cell-to-cell communication system of *Pseudomonas aeruginosa*. Proc Natl Acad Sci U S A. 1999 ; 96(20):11229-11234.

Xie Y, Shi Y H, Wang L L, et al. Outer membrane vesicle contributes to the *Pseudomonas aeruginosa* resistance to antimicrobial peptides in the acidic airway of bronchiectasis patients. MedComm.2025, 6(2): e70084.

Diggle SP, et al. The *Pseudomonas aeruginosa* quinolone signal molecule overcomes the cell density-dependency of the quorum sensing hierarchy, regulates rhl-dependent genes at the onset of stationary phase and can be produced in the absence of LasR. Mol Microbiol. 2003 ; 50(1):29-43.

Bredenbruch F, et al. The *Pseudomonas aeruginosa* quinolone signal (PQS) has an iron-chelating activity. Environ Microbiol. 2006 ; 8(8):1318-1329.

Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins

This product is for Research Use Only· Not for Human or Veterinary or Therapeutic Use

Tel:781-999-4286 E_email:info@targetmol.com Address:34 Washington Street,Wellesley Hills,MA 02481