

Poncirin

Chemical Properties


CAS No. : 14941-08-3

Formula: C₂₈H₃₄O₁₄

Molecular Weight: 594.56

Appearance: no data available

Storage: Powder: -20°C for 3 years | In solvent: -80°C for 1 year



Biological Description

Description	1. Poncirin (Isosakuranetin-7-neohesperidoside) shows a significant in vitro inhibitory effect on the growth of the human gastric cancer cells, SGC-791, in a dose-dependent manner. 2. Poncirin prevents adipogenesis, enhances osteoblast differentiation in mesenchymal stem cells, increased bone mineral density, and improves trabecular microarchitecture likely reflect increases bone formation and decreases bone resorption in GIO mice.
Targets(IC50)	Apoptosis

Solubility Information

Solubility	Ethanol: Soluble, Pyridine, Methanol, etc.: Soluble, DMSO: 91 mg/mL (153.05 mM), Sonication is recommended. (< 1 mg/ml refers to the product slightly soluble or insoluble)
------------	--

Preparing Stock Solutions

	1mg	5mg	10mg
1 mM	1.6819 mL	8.4096 mL	16.8192 mL
5 mM	0.3364 mL	1.6819 mL	3.3638 mL
10 mM	0.1682 mL	0.841 mL	1.6819 mL
50 mM	0.0336 mL	0.1682 mL	0.3364 mL

Please select the appropriate solvent to prepare the stock solution, according to the solubility of the product in different solvents. Please use it as soon as possible.

Reference

Kim D H , Bae E A , Han M J . Anti-Helicobacter pylori activity of the metabolites of poncirin from Poncirus trifoliata by human intestinal bacteria.[J]. Biological & Pharmaceutical Bulletin, 1999, 22(4):422.
Sakaki, Mika, et al. Medicine and food with particular reference to chinpi, dried citrus peel, and a component of Ninjin'yoeito. Neuropeptides. (2021): 102166.

Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins

This product is for Research Use Only· Not for Human or Veterinary or Therapeutic Use

Tel:781-999-4286 E_mail:info@targetmol.com Address:36 Washington Street,Wellesley Hills,MA 02481