Data Sheet (Cat.No.T4008)



Metergoline

Chemical Properties

CAS No.: 17692-51-2


Formula: C25H29N3O2

Molecular Weight: 403.52

Appearance: no data available

store at low temperature

Storage: Powder: -20°C for 3 years | In solvent: -80°C for 1 year



Biological Description

Description	Metergoline (Methergoline) is a dopamine agonist and serotonin antagonist. It has been used similarly to BROMOCRIPTINE as a dopamine agonist and also for migraine disorder therapy.
Targets(IC50)	5-HT Receptor,Dopamine Receptor,Sodium Channel

Solubility Information

Solubility	DMSO: 40.4 mg/mL (100.12 mM), Sonication is recommended.
	(< 1 mg/ml refers to the product slightly soluble or insoluble)

Preparing Stock Solutions

	1mg	5mg	10mg	
1 mM	2.4782 mL	12.391 mL	24.7819 mL	
5 mM	0.4956 mL	2.4782 mL	4.9564 mL	
10 mM	0.2478 mL	1.2391 mL	2.4782 mL	
50 mM	0.0496 mL	0.2478 mL	0.4956 mL	

Please select the appropriate solvent to prepare the stock solution, according to the solubility of the product in different solvents. Please use it as soon as possible.

Page 1 of 2 www.targetmol.com

Reference

Millan, M.J., Newman-Tancredi, A., Lochon, S., et al. Specific labelling of serotonin 5-HT1B receptors in rat frontal cortex with the novel, phenylpiperazine derivative, [3H]GR125,743. A pharmacological characterization. Pharmacol. Biochem. Behav. 71(4), 589-598 (2002).

Knight, A.R., Misra, A., Quirk, K., et al. Pharmacological characterisation of the agonist radioligand binding site of 5-HT2A, 5-HT2B and 5-HT2C receptors. Naunyn Schmiedebergs Arch. Pharmacol. 370(2), 114-123 (2004)

Knight, J.A., Smith, C., Toohey, N., et al. Pharmacological analysis of the novel, rapid, and potent inactivation of the human 5-Hydroxytryptamine7 receptor by risperidone, 9-OH-Risperidone, and other inactivating antagonists. Mol. Pharmacol. 75(2), 374-380 (2009).

Yeom, H.D. and Lee, J.-H. Regulation of human Kv1.4 channel activity by the antidepressant metergoline. Biol. Pharm. Bull. 39(6), 1069-1072 (2016).

Lee, J.-H., Liu, J., Shin, M., et al. Metergoline inhibits the neuronal Nav1.2 voltage-dependent Na+ channels expressed in Xenopus oocytes. Acta. Pharmacol. Sin. 35(7), 862-868 (2014).

Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins

This product is for Research Use Only · Not for Human or Veterinary or Therapeutic Use

Tel:781-999-4286 E_mail:info@targetmol.com Address:36 Washington Street,Wellesley Hills,MA 02481

Page 2 of 2 www.targetmol.com