


Neoline

Chemical Properties

CAS No. :	466-26-2
Formula:	C ₂₄ H ₃₉ NO ₆
Molecular Weight:	437.57
Appearance:	no data available
Storage:	keep away from direct sunlight
	Powder: -20°C for 3 years In solvent: -80°C for 1 year



Biological Description

Description	Neoline is an active ingredient of <i>A. venetum</i> root (PA) that ameliorates oxaliplatin-induced peripheral neuropathy in mice. Neoline inhibits Nav1.7 voltage-gated sodium channel currents (VGSC). It can be used as a labeling compound to determine the quality of PA products used in the study of neuropathic pain.
Targets(IC50)	Sodium Channel

Solubility Information

Solubility	Chloroform, Dichloromethane, Ethyl Acetate, Acetone, etc.: Soluble, DMSO: 50 mg/mL (114.27 mM), Sonication is recommended. (< 1 mg/ml refers to the product slightly soluble or insoluble)
------------	---

Preparing Stock Solutions

	1mg	5mg	10mg
1 mM	2.2853 mL	11.4267 mL	22.8535 mL
5 mM	0.4571 mL	2.2853 mL	4.5707 mL
10 mM	0.2285 mL	1.1427 mL	2.2853 mL
50 mM	0.0457 mL	0.2285 mL	0.4571 mL

Please select the appropriate solvent to prepare the stock solution, according to the solubility of the product in different solvents. Please use it as soon as possible.

Reference

Suzuki, T., Miyamoto, K., Yokoyama, N., Sugi, M., Kagioka, A., & Kitao, Y. et al. (2016). Processed aconite root and its active ingredient neoline may alleviate oxaliplatin-induced peripheral neuropathic pain. *Journal Of Ethnopharmacology*, 186, 44-52. doi: 10.1016/j.jep.2016.03.056

Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins

This product is for Research Use Only. Not for Human or Veterinary or Therapeutic Use

Tel: 781-999-4286 E_mail: info@targetmol.com Address: 36 Washington Street, Wellesley Hills, MA 02481