

URMC-099

## Chemical Properties

CAS No. : 1229582-33-5


Formula: C<sub>27</sub>H<sub>27</sub>N<sub>5</sub>

Molecular Weight: 421.54

Appearance: no data available

Storage: store at low temperature

Powder: -20°C for 3 years | In solvent: -80°C for 1 year



## Biological Description

Description	URMC-099 is an orally bioavailable, brain penetrant MLK inhibitor (IC <sub>50</sub> : 19/42/14/150 nM, for MLK1/MLK2/MLK3/DLK), and also inhibits LRRK2 activity (IC <sub>50</sub> : 11 nM).
Targets(IC <sub>50</sub> )	MLK,MEK,Syk,Bcr-Abl,CDK,Aurora Kinase,Autophagy,c-Met/HGFR,DNA Alkylation,IGF-1R,LRRK2,ROCK,SGK,Src,Trk receptor,VEGFR
In vitro	URMC-099, administered intraperitoneally (i.p.) at a dosage of 10 mg/kg, demonstrates the ability to reduce the production of inflammatory cytokines within the body, protect neuronal structures, and alter the morphology and ultrastructural response of microglia to exposure to HIV-1 Tat. In mice, URMC-099 exhibits favorable pharmacokinetics and enhanced Central Nervous System (CNS) penetration. Additionally, URMC-099 significantly reduces the infiltration of neutrophils into the peritoneum of wild-type mice in response to fMLP.
In vivo	URMC-099 reduces chemotaxis in wild-type neutrophils induced by fMLP in vitro. Moreover, URMC-099 inhibits the release of TNFα in microglia triggered by lipopolysaccharides, as well as the release of cytokines in human monocytes stimulated by HIV-1 Tat. Additionally, URMC-099 prevents the destruction and phagocytosis of cultured neuronal axons by microglia.
Kinase Assay	HDAC IC <sub>50</sub> Profiling: The in vitro inhibitory activity of compounds against seven human HDAC isoforms (1, 2, 4 C2A, 5 C2A, 6, 8, and 11) are performed with a fluorescent based assay according to the company's standard operating procedure. The IC <sub>50</sub> values are determined using 10 different concentrations with 3-fold serial dilution starting at 10 μM. TSA and vorinostat are used as reference compounds.

## Solubility Information

Solubility	H <sub>2</sub> O: < 1 mg/mL (insoluble or slightly soluble), Ethanol: 6 mg/mL (14.23 mM),Sonication is recommended. DMSO: 75 mg/mL (177.92 mM),Sonication is recommended. (< 1 mg/ml refers to the product slightly soluble or insoluble)
------------	--

### Preparing Stock Solutions

	1mg	5mg	10mg
1 mM	2.3723 mL	11.8613 mL	23.7225 mL
5 mM	0.4745 mL	2.3723 mL	4.7445 mL
10 mM	0.2372 mL	1.1861 mL	2.3723 mL
50 mM	0.0474 mL	0.2372 mL	0.4745 mL

Please select the appropriate solvent to prepare the stock solution, according to the solubility of the product in different solvents. Please use it as soon as possible.

### Reference

Goodfellow VS, et al. J Med Chem. 2013, 56(20), 8032-8048.

Marker DF, et al. J Neurosci. 2013, 33(24), 91998-1202010.

Polesskaya O, et al. Mol Immunol. 2014, 58(2), 214-222.

**Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins**

**This product is for Research Use Only. Not for Human or Veterinary or Therapeutic Use**

Tel:781-999-4286    E\_mail:info@targetmol.com    Address:36 Washington Street,Wellesley Hills,MA 02481