


Dovitinib Dilactic Acid

Chemical Properties

CAS No. : 852433-84-2
 Formula: $C_{21}H_{21}FN_6O \cdot 2C_3H_6O_3$
 Molecular Weight: 572.59
 Appearance: no data available
 Storage: Powder: -20°C for 3 years | In solvent: -80°C for 1 year



Biological Description

Description	Dovitinib Dilactic Acid (Dovitinib (TKI-258) Dilactic Acid) is the Dilactic acid of Dovitinib, which is a multitargeted RTK inhibitor, mostly for class III (FLT3/c-Kit) with IC ₅₀ of 1 nM/2 nM, also potent to class IV (FGFR1/3) and class V (VEGFR1-4) RTKs with IC ₅₀ of 8-13 nM, less potent to InsR, EGFR, c-Met, EphA2, Tie2, IGF1R and HER2. Phase 4.
Targets(IC ₅₀)	FGFR,FLT,c-Kit,VEGFR
In vitro	Dovitinib potently inhibits the FGF-stimulated growth of WT and F384L-FGFR3-expressing B9 cells with IC ₅₀ of 25 nM. In addition, Dovitinib inhibits proliferation of B9 cells expressing each of the various activated mutants of FGFR3. Interestingly, there are minimal observed differences in the sensitivity of the different FGFR3 mutations to Dovitinib, with the IC ₅₀ ranging from 70 to 90 nM for each of the various mutations. IL-6-dependent B9 cells containing vector only (B9-MINV cells are resistant to the inhibitory activity of Dovitinib at concentrations up to 1 μM. Dovitinib inhibits cell proliferation of KMS11 (FGFR3-Y373C), OPM2 (FGFR3-K650E), and KMS18 (FGFR3-G384D) cells with IC ₅₀ of 90 nM (KMS11 and OPM2) and 550 nM, respectively. Dovitinib inhibits FGF-mediated ERK1/2 phosphorylation and induces cytotoxicity in FGFR3-expressing primary MM cells. BMSCs does confer a modest degree of resistance with 44.6% growth inhibition for cells treated with 500 nM Dovitinib and cultured on stroma compared with 71.6% growth inhibition for cells grown without BMSCs. Dovitinib inhibits proliferation of M-NFS-60, an M-CSF growth-driven mouse myeloblastic cell line with a median effective concentration (EC ₅₀) of 220 nM. [1] Treatment of SK-HEP1 cells with Dovitinib results in a dose-dependent reduction in cell number and G2/M phase arrest with reduction in the G0/G1 and S phases, inhibition of anchorage-independent growth and blockage of bFGF-induced cell motility. The IC ₅₀ for Dovitinib in SK-HEP1 cells is approximately 1.7 μM. Dovitinib also significantly reduces the basal phosphorylation levels of FGFR-1, FGFR substrate 2α (FRS2-α) and ERK1/2 but not Akt in both SK-HEP1 and 21-0208 cells. In 21-0208 HCC cells, Dovitinib significantly inhibits bFGF-induced phosphorylation of FGFR-1, FRS2-α, ERK1/2 but not Akt. [2]
In vivo	Dovitinib induces both cytostatic and cytotoxic responses in vivo resulting in regression of FGFR3-expressing tumors.[1] Dovitinib shows a dose- and exposure-dependent inhibition of target receptor tyrosine kinases (RTKs) expressed in tumor xenografts. Dovitinib potently inhibits tumor growth of six HCC lines. Inhibition of angiogenesis correlated with inactivation of FGFR/PDGFRβ/VEGFR2 signaling pathways. In an orthotopic model, Dovitinib potently inhibits primary tumor growth and lung metastasis and significantly prolonged mouse survival. [2] Administration of Dovitinib results in

	significant tumor growth inhibition and tumor regressions, including large, established tumors (500-1,000 mm ³). [3]
Kinase Assay	In vitro kinase assays: The inhibitory concentration of 50% (IC ₅₀) values for the inhibition of RTKs by Dovitinib are determined in a time-resolved fluorescence (TRF) or radioactive format, measuring the inhibition by Dovitinib of phosphate transfer to a substrate by the respective enzyme. The kinase domains of FGFR3, FGFR1, PDGFR β , and VEGFR1-3 are assayed in 50 mM HEPES (N-2-hydroxyethylpiperazine-N'-2-ethanesulfonic acid), pH 7.0, 2 mM MgCl ₂ , 10 mM MnCl ₂ 1 mM NaF, 1 mM dithiothreitol (DTT), 1 mg/mL bovine serum albumin (BSA), 0.25 μ M biotinylated peptide substrate (GGGGQDGKDYIVLPI), and 1 to 30 μ M adenosine triphosphate (ATP) depending on the K _m for the respective enzyme. ATP concentrations are at or just below K _m . For c-KIT and FLT3 reactions the pH is raised to 7.5 with 0.2 to 8 μ M ATP in the presence of 0.25 to 1 μ M biotinylated peptide substrate (GGLFDDPSYVNVQNL). Reactions are incubated at room temperature for 1 to 4 hours and the phosphorylated peptide captured on streptavidin-coated microtiter plates containing stop reaction buffer (25 mM EDTA [ethylenediaminetetraacetic acid], 50 mM HEPES, pH 7.5). Phosphorylated peptide is measured with the DELFIA TRF system using a Europium-labeled antiphosphotyrosine antibody PT66. The concentration of Dovitinib for IC ₅₀ is calculated using nonlinear regression with XL-Fit data analysis software version 4.1 (IDBS). Inhibition of colony-stimulating factor-1 receptor (CSF-1R), PDGFR α , insulin receptor (InsR), and insulin-like growth factor receptor 1 (IGFR1) kinase activity is determined at ATP concentrations close the K _m for ATP.
Cell Research	Cell viability is assessed by 3-(4,5-dimethylthiazol)-2,5-diphenyl tetrazolium (MTT) dye absorbance. Cells are seeded in 96-well plates at a density of 5 $\times$ 10 ³ (B9 cells) or 2 $\times$ 10 ⁴ (MM cell lines) cells per well. Cells are incubated with 30 ng/mL aFGF and 100 μ g/mL heparin or 1% IL-6 where indicated and increasing concentrations of Dovitinib. For each concentration of Dovitinib, 10 μ L aliquots of drug or DMSO diluted in culture medium is added. For drug combination studies, cells are incubated with 0.5 μ M dexamethasone, 100 nM Dovitinib, or both simultaneously where indicated. To evaluate the effect of Dovitinib on growth of MM cells adherent to BMSCs, 104 KMS11 cells are cultured on BMSC-coated 96-well plates in the presence or absence of Dovitinib. Plates are incubated for 48 to 96 hours. For assessment of macrophage colony-stimulating factor (M-CSF)-mediated growth, 5 $\times$ 10 ³ M-NFS-60 cells/well are incubated with serial dilutions of Dovitinib with 10 ng/mL M-CSF and without granulocyte-macrophage colony-stimulating factor (GM-CSF). After 72 hours cell viability is determined using Cell Titer-Glo Assay. Each experimental condition is performed in triplicate. (Only for Reference)

Solubility Information

Solubility	DMSO: 83 mg/mL (144.96 mM), Sonication is recommended. H ₂ O: 64 mg/mL (111.77 mM), Sonication is recommended. Ethanol: <1 mg/mL, (< 1 mg/mL refers to the product slightly soluble or insoluble)
------------	---

Preparing Stock Solutions

	1mg	5mg	10mg
1 mM	1.7465 mL	8.7323 mL	17.4645 mL
5 mM	0.3493 mL	1.7465 mL	3.4929 mL
10 mM	0.1746 mL	0.8732 mL	1.7465 mL
50 mM	0.0349 mL	0.1746 mL	0.3493 mL

Please select the appropriate solvent to prepare the stock solution, according to the solubility of the product in different solvents. Please use it as soon as possible.

Reference

Trudel S, et al. Blood, 2005, 105(7), 2941-2948.

Huynh H, et al. J Hepatol. 2012, 56(3), 595-601.

Lee SH, et al. Clin Cancer Res. 2005, 11(10), 3633-3641.

Azab AK, et al. Clin Cancer Res, 2011, 17(13), 4389-4399.

Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins

This product is for Research Use Only· Not for Human or Veterinary or Therapeutic Use

Tel:781-999-4286 E_mail:info@targetmol.com Address:36 Washington Street,Wellesley Hills,MA 02481