


## Vilazodone Hydrochloride

### Chemical Properties

CAS No. :	163521-08-2
Formula:	C <sub>26</sub> H <sub>27</sub> N <sub>5</sub> O <sub>2</sub> ·HCl
Molecular Weight:	477.99
Appearance:	no data available
Storage:	Powder: -20°C for 3 years   In solvent: -80°C for 1 year



### Biological Description

Description	Vilazodone Hydrochloride (EMD 68843) , a partial agonist of 5-HT <sub>1A</sub> receptors and specific serotonin reuptake inhibitor (SSRI), is utilized in treating of the major depressive disorder.
Targets(IC <sub>50</sub> )	5-HT Receptor,Serotonin Transporter
In vitro	Vilazodone demonstrates an IC <sub>50</sub> of 0.2 nM at the human 5-HT <sub>1A</sub> receptor and 0.5 nM for the SERT. Vilazodone displays high affinity (pK <sub>i</sub> ≥ 9.3) for human recombinant and rat, guinea pig, mouse, and marmoset native tissue 5-HT <sub>1A</sub> receptors. [1]
In vivo	Vilazodone selectively enhances serotonergic output in the prefrontal cortex of rats. Vilazodone demonstrates anxiolytic efficacy through Behavioral evaluations in the ultrasonic vocalization model of anxiety in rats. Vilazodone also shows efficacy but at a single dose only in the forced swim test (a putative model of depression). [1] Vilazodone (1-10 mg/kg p.o.) dose-dependently displaces in vivo [ <sup>3</sup> H]DASB (N,N-dimethyl-2-(2-amino-4-cyanophenylthio)benzylamine) binding from rat cortex and hippocampus, indicating that vilazodone occupies 5-HT transporters in vivo. Vilazodone (10 mg/kg p. o.) is demonstrated to cause a 2-fold increase in extracellular 5-HT but no change in noradrenaline or dopamine levels in frontal cortex of freely moving rats. [2] Vilazodone affects stress potentiation of startle at doses above 5 mg/kg in rats. Vilazodone increases stress elevation of startle at 10 mg/kg in rats. Vilazodone (20 and 40 mg/kg) blocked stress potentiation of startle in rats. Vilazodone increases stress elevation of startle at all doses in rats. [3] vilazodone has no effect on 5-HT efflux at 100 nM but significantly decreases 5-HT efflux at 1 mM in the guinea-pig dorsal raphe nucleus. Vilazodone significantly increases the re-uptake half life for 5-HT in the guinea-pig dorsal raphe nucleus. [4]

### Solubility Information

Solubility	DMSO: 45 mg/mL (94.14 mM),Sonication is recommended. (< 1 mg/ml refers to the product slightly soluble or insoluble)
------------	---

### Preparing Stock Solutions

	1mg	5mg	10mg
1 mM	2.0921 mL	10.4605 mL	20.9209 mL
5 mM	0.4184 mL	2.0921 mL	4.1842 mL
10 mM	0.2092 mL	1.046 mL	2.0921 mL
50 mM	0.0418 mL	0.2092 mL	0.4184 mL

Please select the appropriate solvent to prepare the stock solution, according to the solubility of the product in different solvents. Please use it as soon as possible.

### Reference

Dawson LA, et al. CNS Neurosci Ther, 2009, 15(2), 107-117.

Hughes ZA, et al. Eur J Pharmacol, 2005, 510(1-2), 49-57.

Adamec R, et al. Eur J Pharmacol, 2004, 504(1-2), 65-77.

**Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins**

**This product is for Research Use Only. Not for Human or Veterinary or Therapeutic Use**

Tel:781-999-4286 E\_mail:info@targetmol.com Address:36 Washington Street,Wellesley Hills,MA 02481