


## Embelin

## Chemical Properties

CAS No. :	550-24-3
Formula:	C <sub>17</sub> H <sub>26</sub> O <sub>4</sub>
Molecular Weight:	294.39
Appearance:	no data available
Storage:	keep away from direct sunlight Powder: -20°C for 3 years   In solvent: -80°C for 1 year



## Biological Description

Description	Embelin (Embelic acid), isolated from the Japanese Ardisia herb, is an inhibitor of the X-linked inhibitor of apoptosis (IC <sub>50</sub> : 4.1 μM).
Targets(IC <sub>50</sub> )	Apoptosis,NF-κB,Autophagy,IAP,Lipoxygenase,Prostaglandin Receptor
In vitro	Embelin from the Japanese Ardisia herb is a small-molecular inhibitor that binds to the XIAP BIR3 domain with which Smac and caspase-9 bind. Embelin inhibits cell growth of both PC-3 and LNCap cells in a dose-dependent manner, with IC <sub>50</sub> values of 3.7 and 5.7 μM. While, the toxicity of Embelin in normal PrEC and in WI-38 cells is much lower with IC <sub>50</sub> values of 20.1 μM and 19.3 μM. Treatment of PC-3 cells with 25 and 50 μM of embelin for 48 h, 30% and 75% of cells undergo apoptosis, representing approximately 3- and 9-fold increase as compared to untreated cells. [1] Embelin also potently suppresses the biosynthesis of eicosanoids by selective inhibition of 5-lipoxygenase (5-LO) and Microsomal prostaglandin E <sub>2</sub> synthase-1 (mPGES-1) with IC <sub>50</sub> = 0.06 and 0.2 mM, respectively. [2]
Kinase Assay	Fluorescence Polarization Competitive Binding Assay: Fluorescence polarization experiments are performed in Dynex 96-well, black, round-bottom plates. A 5 μL sample of Embelin dilutions in DMSO, and preincubated XIAP BIR3 protein (0.06 μM) and the N terminus of a Smac peptide (SM7F) (0.01 μM) in the assay buffer are added to 96-well plates to produce a final volume of 125 μL. For each assay, the bound peptide control containing XIAP BIR3 protein and SM7F (equivalent to 0% inhibition) and free peptide control containing only free SM7F (equivalent to 100% inhibition) were included. The plates were mixed and incubated at room temperature for 3 h to reach equilibrium.
Cell Research	Cell growth is determined by the MTT-based assay using Cell Proliferation Reagent WST-1 according to the manufacturer's instruction. Cells (5000 cells/well) are grown in medium with 10% FBS, and various concentrations of Embelin are added to the cells. Four to five days later, WST-1 is added to each well and incubated for 1-3 h at 37 °C. Absorbance is measured with a plate reader at 450 nm with correction at 650 nm.(Only for Reference)

## Solubility Information

## A DRUG SCREENING EXPERT

Solubility	Ethanol: 2.9 mg/mL (10 mM)),Heating is recommended. DMSO: 29.4 mg/mL (99.87 mM),Sonication is recommended. (< 1 mg/ml refers to the product slightly soluble or insoluble)
------------	--

### Preparing Stock Solutions

	1mg	5mg	10mg
1 mM	3.3969 mL	16.9843 mL	33.9685 mL
5 mM	0.6794 mL	3.3969 mL	6.7937 mL
10 mM	0.3397 mL	1.6984 mL	3.3969 mL
50 mM	0.0679 mL	0.3397 mL	0.6794 mL

Please select the appropriate solvent to prepare the stock solution, according to the solubility of the product in different solvents. Please use it as soon as possible.

### Reference

Nikolovska-Coleska Z, et al. J Med Chem, 2004, 47(10), 2430-2440.

Peng C, Wu F, Ma Y, et al.Ginkgolic acid Inhibits Ebola Virus Transcription and Replication by Disrupting the Interaction Between Nucleoprotein and VP30 Protein.Antiviral Research.2024: 106074.

Schaible AM, et al. Biochem Pharmacol, 2013, 86(4), 476-486.

**Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins**

**This product is for Research Use Only· Not for Human or Veterinary or Therapeutic Use**

Tel:781-999-4286 E\_mail:info@targetmol.com Address:36 Washington Street,Wellesley Hills,MA 02481